

Linear systems on graphs with a real structure

Marc Coppens*

Abstract

A degeneration of a smooth projective curve to a strongly stable curve gives rise to a specialization map from divisors on curves to divisors on graphs. In this paper we show that this specialization behaves well under the presence of real structures. In particular we study real linear systems on graphs with a real structure and we prove results on them comparable to results in the classical theory of real curves. We also consider generalizations to metric graphs and tropical curves

1 Introduction

In [1] one introduces a theory of linear systems on graphs which is very similar to the theory of linear systems on smooth complex projective curves. In [2] one shows a relation by means of specialization of a smooth projective curve to a strongly semistable curve. Associated to this strongly semistable curve there is a dual graph G and divisors on the geometric generic smooth fiber give rise to divisors on the metric graph T associated to G . This specialization preserves linear equivalence on curves and on graphs. It is the aim of this paper to show that this specialization is compatible with real structures. This means in case there is a real structure on the family of curves there is a real structure on the graph and on the metric graph and real divisors on curves specialize to real divisors on the graph.

In Section 2 we introduce the notion of a graph G with a real structure (Definition 2.1). We define an associated real locus graph $G(\mathbb{R})$ (Definition 2.5). This is similar to the notion of the real locus $X(\mathbb{R})$ of a real curve X . In this classical case there are strong constraints on the number of connected components $s(X)$ of $X(\mathbb{R})$, the genus $g(X)$ and a number $a(X)$ determined by the number of connected components of $X(\mathbb{C}) \setminus X(\mathbb{R})$. A graph G has a genus $g(G)$ and we define similar numbers $s(G)$ (Formula 2.1) and $a(G)$ (Definition 2.6). The definition of $s(G)$ not only takes into account the number of connected components of $G(\mathbb{R})$ but also their genus (in the curve case all components of $X(\mathbb{R})$ are circles). In the graph case we prove there are similar but stronger

*Katholieke Hogeschool Kempen, Departement IBW, Kleinhoefstraat 4, B-2440 Geel, Belgium; K.U.Leuven, Departement of Mathematics, Celestijnenlaan 200B, B-3001 Leuven, Belgium; email: marc.coppens@khk.be

constraints (Theorem 2.7). We prove that all cases satisfying those constraints do occur (Example 2.8).

As mentioned before, there is a theory of divisors and linear systems on graphs similar to the theory on complex curves. In case the graph has a real structure one can consider real divisors and real linear systems. In Section 3 we prove results similar to basic properties of real linear systems on real curves. In particular we prove that the parity of the degree of the restriction of a real divisor D to a connected component of $G(\mathbb{R})$ does not change under linear equivalence (Theorem 3.5). We introduce an M-graph as being a graph satisfying $s(G) = g(G) + 1$ (Definition 3.7; this is similar to the concept of an M-curve). We prove that each real effective divisor on an M-graph is linearly equivalent to a totally real effective divisor (Theorem 3.9). A similar strong fact does not hold in the case of M-curves. In case $G(\mathbb{R})$ has $g(G) + 1$ connected components we say that G is a strong M-graph (Definition 3.10) and we prove G has a real linear system g_2^1 in that case (Proposition 3.11).

In order to study the compatibility of the above mentioned specialization with real structures we need to consider metric graphs with a real structure. This is done in Section 4. We generalize the results on real linear systems on graphs to the context of metric graphs with a real structure. We also make further generalizations to tropical curves.

Finally in Section 5 we study the behavior of the real structure under specialization. Although at the end this follows the arguments of [2] quite closely (see the proof of Theorem 5.8), we need to take care about this behavior of the real structure in those arguments. In particular for the generic fiber of the degeneration the base field is a real field and we use the real closure instead of the algebraic closure to define real divisors on the generic fiber. Therefore we need to pay some attention to extensions of real fields in those arguments (Lemmas 5.1, 5.2 and 5.3). Also we need to take care about the existence of real structures on certain desingularizations occurring in those arguments (Lemma 5.5) and we need to check that this real structure is compatible to the real structure on the metric graph T associated to G (Proposition 5.7). Finally we prove that all graphs with a real structure can be obtained from such a degeneration (Proposition 5.9).

2 Graphs with a real structure

We recall some generalities on finite graphs (as a general reference, see e.g. [4]). A *finite graph* G is defined by a finite set of vertices $V(G)$ and a finite set of edges $E(G)$ together with an incidence function that associates to an edge e a subset $\psi(e) \subset V(G)$ containing one or two vertices. In case it contains only one vertex v we say e is a *loop* at v . When we write $\psi(e) = \{v, w\}$ it is possible that $v = w$ (i.e. e could be a loop). Vertices of $\psi(e)$ are called the *ends* of e . We write $v(G)$ (resp. $e(G)$) to denote the number of vertices (resp. edges) of G .

A *walk* Γ in G is a sequence $v_0 e_0 v_1 e_1 v_2 \cdots v_{n-1} e_{n-1} v_n$ such that $e_i \in E(G)$ with $\psi(e_i) = \{v_i, v_{i+1}\}$ for $0 \leq i \leq n-1$. We call v_0 and v_n the *ends* of the

walk Γ and we say Γ connects v_0 to v_n . We call e_i ($0 \leq i \leq n-1$) the edges of Γ and v_i ($1 \leq i \leq n-1$) the internal vertices of Γ . A walk Γ is called closed if the ends v_0 and v_n of Γ are equal. A graph G is *connected* if for each two different vertices v_1, v_2 of G there exists a walk having ends v_1 and v_2 .

A *subgraph* G' of G is defined by subsets $V(G') \subset V(G)$ and $E(G') \subset E(G)$ such that for $e \in E(G')$ one has $\psi(e) \subset V(G')$. If G' and G'' are two subgraphs of G then $G' \cup G''$ is the subgraph defined by $V(G') \cup V(G'') \subset V(G)$ and $E(G') \cup E(G'') \subset E(G)$. A subgraph C of G is called a *cycle* if all vertices and edges of C are the vertices and edges of a closed walk such that the end vertices are the only vertices being equal in that walk. A *connected component* of a graph G is a subgraph G' such that, each subgraph $G'' \neq G'$ containing G' is not connected. A graph G has finitely many connected components $G_1; \dots; G_c$. We write $c = c(G)$ to denote the number of connected components of G . The *genus* of the graph G is defined by

$$g(G) = c(G) + e(G) - v(G).$$

From now on in this paper G denotes a connected graph.

Definition 2.1. Let G be a finite graph. A *real structure* on G is defined by an involution $\iota : G \rightarrow G$ (hence $\iota \circ \iota$ is the identity on G). More concretely, it is defined by involutions $\iota_V : V(G) \rightarrow V(G)$ and $\iota_E : E(G) \rightarrow E(G)$ such that for each $e \in E(G)$ one has $\psi(\iota_E(e)) = \iota_V(\psi(e))$.

From now on we assume G is a graph with a fixed real structure. For $v \in V(G)$ (resp. $e \in E(G)$) we write $\bar{v}$ (resp. $\bar{e}$) instead of $\iota_V(v)$ (resp. $\iota_E(e)$).

Definition 2.2. A vertex $v \in V(G)$ is called a *real vertex* if $v = \bar{v}$. In case v is not a real vertex $\bar{v}$ is called the *conjugated vertex*. In that case the symbol $v + \bar{v}$ (a real divisor on G according to the terminology of Section 3) is called a *non-real vertex pair* of G .

We write $V_{\mathbb{R}}(G)$ to denote the subset of $V(G)$ of real vertices of G .

Definition 2.3. An edge $e \in E(G)$ is called a *real edge* if $e = \bar{e}$. A real edge e is called an *isolated real edge* if $\psi(e) \not\subset V_{\mathbb{R}}(G)$, otherwise we say e is a *non-isolated real edge*. In case e is not a real edge $\bar{e}$ is called the *conjugated edge* of e .

We write $E_{\mathbb{R}}(G)$ (resp. $E_{\mathbb{R}}^0(G)$) to denote the subset of $E(G)$ of real edges (resp. non-isolated real edges) of G .

Definition 2.4. A subgraph G' of G is called a *real subgraph* if the subsets $V(G')$ of $V(G)$ and $E(G')$ of $E(G)$ are invariant under ι_V and ι_E .

If G' is a real subgraph of G then the restrictions of ι_V to $V(G')$ and ι_E to $E(G')$ induce a real structure on G' . We always consider this real structure on G' , hence a real subgraph of G is always considered as a graph with a real structure.

Definition 2.5. *The subgraph of G having $V_{\mathbb{R}}(G)$ (resp. $E_{\mathbb{R}}^0(G)$) as its vertex set (resp. edge set) is called the real locus graph of G . It is denoted by $G(\mathbb{R})$.*

Clearly $G(\mathbb{R})$ is a real subgraph of G and the identity defines the real structure on $G(\mathbb{R})$. The number of connected components of $G(\mathbb{R})$ is denoted by $s'(G)$. We denote the components by $G(\mathbb{R})_i$ ($1 \leq i \leq s'(G)$). Let $e^i(G)$ be the number of isolated real edges of G . Now we introduce the number $s(G)$ that corresponds to the number of connected components of $X(\mathbb{R})$ in case X is a real curve as follows:

$$s(G) = e^i(G) + \sum_{i=1}^{s'(G)} (g(G(\mathbb{R})_i) + 1). \quad (2.1)$$

Remark. The definition of this number $s(G)$ is in accordance with the degeneration studied in Section 5. Consider the graph G as the dual graph of a total degeneration of a smooth real curve of genus $g(G)$ (see Proposition 5.9). In such degeneration one should compare $G(\mathbb{R})_i$ to a real algebraic curve of genus $g(G(\mathbb{R})_i)$, the real locus of such curve has at most $g(G(\mathbb{R})_i) + 1$ connected components. On the other hand, an isolated real edge corresponds to an isolated real singular point of a real curve. Such isolated real singular point can deform to a connected component of the real locus of a smooth curve. In this way $s(G)$ corresponds to the maximal number of connected components of the real locus of a smooth real curve having a total degeneration with dual graph G with its real structure.

Next we define the number $a(G)$ similar to the number $a(X)$ of a real curve X defined by $a(X) = 1$ if $X(\mathbb{C}) \setminus X(\mathbb{R})$ is connected and $a(X) = 0$ otherwise.

Definition 2.6. *Let G be a graph with a real structure. In case there exists a non-real vertex v and a walk Γ in G with ends v and $\bar{v}$ such that Γ contains neither a real vertex, nor a real edge, we put $a(G) = 1$. Otherwise we put $a(G) = 0$.*

The following theorem gives conditions between the integers $g(G)$, $a(G)$ and $s(G)$.

Theorem 2.7. *Let G be a graph with a real structure. One has $s(G) \equiv g(G) + 1 \pmod{2}$ and $0 \leq s(G) \leq g(G) + 1$. Moreover $s(G) \neq g(G) + 1$ in case $a(G) = 1$ and $s(G) \neq 0$ in case $a(G) = 0$.*

Remark. In the case of a real smooth curve X the condition $s(X) \equiv g(X) + 1 \pmod{2}$ need not hold if $a(X) = 1$. Of course, in the case of graphs, this condition implies $s(G) \leq g(G) - 1$ in case $a(G) = 1$.

Before giving the proof of Theorem 2.7 we introduce some notations we use in that proof. Let G be a graph and let W be a subset of $V(G)$. Then $G[W]$ is the subgraph induced by W . This means, the set of vertices of $G[W]$ is equal to W and the edges of $G[W]$ are the edges e of G such that $\psi(e) \subset W$. Let S be a subset of $E(G)$ then $G[S]$ is the subgraph of G whose set of edges is equal

to S and the set of vertices of $G[S]$ is the union $\bigcup_{e \in S} \psi(e)$. The graph $G \setminus S$ is the subgraph of G obtained by deleting the edges belonging to S (hence its set of vertices is equal to $V(G)$). Finally $G(S)$ is the graph obtained from G by contracting each edge not contained in S . Since G is connected it is obtained from $G[S]$ by identifying two vertices v and v' if and only if there is a walk P in $G \setminus S$ having ends v and v' . For a subset S of $E(G)$ we use the equation

$$g(G) = g(G(S)) + g(G \setminus S). \quad (2.2)$$

Clearly, even if G is a connected graph, the graph $G \setminus S$ need not be connected. In particular it can have isolated vertices as some connected components. Such components do not contribute to $g(G \setminus S)$.

Proof of Theorem 2.7.

Step 1. Reduction to the case that G has no isolated real edges.

Assume e is an isolated real edge of G with $\psi(e) = \{v; \bar{v}\}$ for some non-real vertex pair $v + \bar{v}$. Let G' be the graph with real structure obtained from G by first deleting the edge e (this is a real subgraph of G) and then adding a new vertex v_e and two new edges e' and e'' with $\psi(e') = \{v; v_e\}$ and $\psi(e'') = \{\bar{v}; v_e\}$ and putting $\iota(v_e) = v_e$ and $\iota(e') = e''$, $\iota(e'') = e'$. Instead of the isolated real edge e this graph has one more component $\{v_e\}$ of $G'(\mathbb{R})$. Clearly $g(G') = g(G)$, $s(G') = s(G)$ and $a(G') = a(G)$. Continuing in the way this shows it is enough to prove the case without isolated real edges. So from now on we assume $e^i(G) = 0$.




Figure 1: Replacing an isolated real edge by a real vertex.

Step 2. The case $G(\mathbb{R})$ is empty.

In case $G(\mathbb{R})$ is empty one has $a(G) = 1$ and $s(G) = 0$ and we need to prove that $g(G)$ is odd. Choose a vertex v of G and a walk Γ in G that connects v to $\bar{v}$. Writing $\Gamma = v_0(=v)e_0v_1 \cdots v_{n-1}e_{n-1}v_n(=\bar{v})$ we can assume that for each $0 \leq i < j \leq n$ we have $v_j \notin \{v_i; \bar{v}_i\}$, otherwise we replace Γ by a shorter walk. It follows that $\Gamma \cup \bar{\Gamma}$ is a cycle C with $\bar{C} = C$. Let e be an edge of C with $v \in \psi(e)$ and consider the graph $G' = G \setminus \{e; \bar{e}\}$. Clearly $C \setminus \{e; \bar{e}\}$ has exactly

two connected components C_1 and C_2 with $C_2 = \overline{C_1}$. Let $w \in V(G) \setminus V(C)$. Since G is connected there exists $w' \in V(C)$ and a walk Γ_w that connects w to w' such that Γ_w contains no edge of C . In case $w' \in V(C_1)$, w belongs to the connected component G_1 of G' containing C_1 . In that case $\overline{\Gamma_w}$ connects $\overline{w}$ to $\overline{w'} \in V(C_2)$, hence $\overline{w}$ belongs to the connected component G_2 of G' containing C_2 .




Figure 2: Making $G(\{e, \bar{e}\})$.

It follows that G' is either connected or it has exactly two connected components G_1 and G_2 . In case G' is not connected $G(\{e, \bar{e}\})$ is a cycle, hence it has genus 1 and $g(G') = 2g(G_1)$. Hence Formula (2.2) implies $g(G) = 1 + 2g(G_1)$, in particular $g(G)$ is odd. In case G' is connected $G(\{e, \bar{e}\})$ consists of one vertex and two loops, hence it has genus 2. Hence Formula (2.2) implies $g(G) \equiv g(G') \pmod{2}$. Moreover G' is a connected graph with a real structure satisfying $s(G') = 0$, hence $a(G') = 0$. Since $\sharp(E(G')) < \sharp(E(G))$ we can use induction and assume $g(G')$ is odd. It follows that $g(G)$ is odd too.

From now on we assume $G(\mathbb{R})$ is not empty.

Step 3. It is enough to prove the theorem in case $a(G) = 0$.

Assume the theorem holds in case $a = 0$ and assume $a(G) = 1$. There exists a non-real vertex pair $v + \bar{v}$ and a walk Γ that connects v to $\bar{v}$ such that Γ contains no real vertex and no real edge. As in the previous step we can assume $C = \Gamma \cup \bar{\Gamma}$ is a cycle and we take e and $\bar{e}$ as in the proof of the previous step. Since $G(\mathbb{R}) \neq \emptyset$ there is some real vertex w . In case $G \setminus \{e, \bar{e}\}$ has two connected components G_1 and G_2 it would follow that $w \in G_1 \cap G_2$ since $G_2 = \overline{G_1}$. This is impossible, hence $G \setminus \{e, \bar{e}\}$ is a connected graph G' with $G'(\mathbb{R}) \neq \emptyset$. As explained in the previous step this graph satisfies $s(G) = s(G')$ and $g(G') = g(G) - 2$. In case $a(G') = 0$, $s(G') \equiv g(G') + 1 \pmod{2}$ and $s(G') \leq g(G') + 1$ implies $s(G) \equiv g(G) + 1 \pmod{2}$ and $s(G) \leq g(G) - 1$. In case $a(G') = 1$ we can continue this procedure that has to finish because the genus becomes less.

From now on we can assume $a(G) = 0$.

Step 4. The case $G(\mathbb{R})$ is connected.

Consider the induced subgraph $G[V_{\mathbb{R}}(G)]$. Of course $G(\mathbb{R})$ is a subgraph of $G[V_{\mathbb{R}}(G)]$ having the same set of vertices $V_{\mathbb{R}}(G)$. If e is an edge in $G[V_{\mathbb{R}}(G)]$ not contained in $G(\mathbb{R})$ then $\psi(e) \subset V_{\mathbb{R}}(G)$, hence $\psi(e) = \psi(\bar{e})$ and therefore $\bar{e}$ is another edge in $G[V_{\mathbb{R}}(G)]$ not contained in $G(\mathbb{R})$. Since $G(\mathbb{R})$ is connected it follows that $g(G[V_{\mathbb{R}}(G)]) \geq g(G(\mathbb{R}))$ with equality if and only if $G[V_{\mathbb{R}}(G)] = G(\mathbb{R})$. Moreover we obtain $g(G[V_{\mathbb{R}}(G)]) \equiv g(G(\mathbb{R})) \pmod{2}$ and since $g(G(\mathbb{R})) = s(G) - 1$ in case $G(\mathbb{R})$ is connected we obtain

$$g(G[V_{\mathbb{R}}(G)]) \equiv s(G) + 1 \pmod{2}. \quad (2.3)$$

This finishes the proof of the theorem in case $G = G[V_{\mathbb{R}}(G)]$, so we assume that $G \neq G[V_{\mathbb{R}}(G)]$.




Figure 3: Example of a graph G with connected $G(\mathbb{R})$.




Figure 4: Graphs G' and $G(E')$ of the example.

Assume $G[V_{\mathbb{R}}(G)]$ is just one real vertex v and contains no edge. Let G' be the graph obtained from G by adding a loop e at v and by extending the real structure by putting $\iota_E(e) = e$. Then $g(G') = g(G) + 1$, $a(G') = a(G)$ and $s(G') = s(G) + 1$. Therefore we can assume $G[V_{\mathbb{R}}(G)]$ is not equal to just one real vertex and no edge. Since $G[V_{\mathbb{R}}(G)]$ is connected it follows that $G[V_{\mathbb{R}}(G)] = G[E']$ with E' the set of edges of $G[V_{\mathbb{R}}(G)]$. Let $G' = G \setminus E'$, it is a real subgraph of G . We are going to use the decomposition (see Formula 2.2)

$$g(G) = g(G(E')) + g(G'). \quad (2.4)$$

The graph $G(E')$ is obtained from $G[E'] = G[V_{\mathbb{R}}(G)]$ by identifying vertices v_1 and v_2 of $V_{\mathbb{R}}(G)$ if and only if there is a walk Γ in G' with ends v_1 and v_2 . In particular we obtain

$$g(G(E')) \geq g(G[V_{\mathbb{R}}(G)]) \geq g(G(\mathbb{R})) = s(G) - 1. \quad (2.5)$$

Together with Formula 2.4 this already proves $s(G) \leq g(G) + 1$. Moreover we obtain the following claim that will be used in Section 3.

Claim. If $G(\mathbb{R})$ is connected then $s(G) = g(G) + 1$ if and only if $G[V_{\mathbb{R}}(G)] = G(\mathbb{R}) = G(E')$ and $g(G') = 0$.

Now we are going to prove $s(G) \equiv g(G) + 1 \pmod{2}$. We write $\overline{V}_{\mathbb{R}}(G)$ to denote the set of vertices of $G(E')$. We know $\overline{V}_{\mathbb{R}}(G)$ can be considered as a quotient set of $V_{\mathbb{R}}(G)$. Clearly one has

$$g(G(E')) = g(G[V_{\mathbb{R}}(G)]) + (\#(V_{\mathbb{R}}(G)) - \#(\overline{V}_{\mathbb{R}}(G))). \quad (2.6)$$

Because of Formula 2.3 this implies

$$g(G(E')) \equiv s(G) + 1 + (\#(V_{\mathbb{R}}(G)) - \#(\overline{V}_{\mathbb{R}}(G))) \pmod{2}. \quad (2.7)$$

We make a partition $V_{\mathbb{R}}(G) = V_{\mathbb{R}}(G)_0 \cup V_{\mathbb{R}}(G)_1$ such that a real vertex v belongs to $V_{\mathbb{R}}(G)_0$ if and only if for each edge e with $v \in \psi(e)$ one has $\psi(e) \subset V_{\mathbb{R}}(G)$. This is equivalent to $\{v\}$ being a component of $G' = G \setminus E'$. We have an induced partition $\overline{V}_{\mathbb{R}}(G) = \overline{V}_{\mathbb{R}}(G)_0 \cup \overline{V}_{\mathbb{R}}(G)_1$ (here $\overline{V}_{\mathbb{R}}(G)_1$ is the quotient set of $V_{\mathbb{R}}(G)_1$ using the identification mentioned before) and we obtain

$$g(G(E')) \equiv s(G) + 1 + (\#(V_{\mathbb{R}}(G)_1) - \#(\overline{V}_{\mathbb{R}}(G)_1)) \pmod{2}. \quad (2.8)$$

Let S' be the set of edges e of G' satisfying $\psi(e)$ contains a real vertex. This set is invariant under ι_E and it contains no real edge, hence $\#(S')$ is even. We consider the decomposition

$$g(G') = g(G'(S')) + g(G' \setminus S'). \quad (2.9)$$

Let $V'_{\mathbb{R}}(G)_1$ be the quotient set of $V_{\mathbb{R}}(G)_1$ obtained by identifying vertices v_1 and v_2 from $V_{\mathbb{R}}(G)_1$ if and only if they are ends of a walk with edges belonging to S' . There is a natural bijection between the set of connected components of




Figure 5: Identification of real vertices of the example.

the graph $G'[S']$ and $V'_R(G)_1$. Moreover $\bar{V}_R(G)_1$ can be considered as a quotient set of $V'_R(G)_1$. Also $\sharp(V(G'[S']) \setminus V_R(G)_1)$ is even. Those remarks imply

$$g(G'[S']) \equiv \sharp(V'_R(G)_1) - \sharp(V_R(G)_1) \pmod{2}. \quad (2.10)$$

The graph $G'(S')$ is obtained from $G'[S']$ by identifying vertices v_1 and v_2 of $G'[S']$ if and only if there is a walk Γ in $G' \setminus S'$ with ends v_1 and v_2 . Since $\psi(e)$ contains no real vertex for an edge e of $G' \setminus S'$ it follows v_1 and v_2 are non-real vertices. Since $a(G) = 0$ also $v_2 \neq \bar{v}_1$. On the other hand, $\bar{\Gamma}$ is a walk in $G' \setminus S'$ with ends $\bar{v}_1$ and $\bar{v}_2$. Hence $\bar{v}_1$ and $\bar{v}_2$ as vertices of $G'[S']$ are also identified in $G'(S')$. This proves $\sharp(V(G'[S'])) - \sharp(V(G'(S')))$ is even.




Figure 6: Graph $G'(S')$ of the example.

Assume C_1 and C_2 are two different connected components of $G'[S']$ represented by v'_1 and v'_2 in $V'_R(G)_1$. Assume C_i has vertex v_i such that v_1 and v_2 are identified in $G'(S')$. This is equivalent to the existence of a walk Γ in $G' \setminus S'$ that connects v_1 to v_2 . Hence it is equivalent to the fact that v'_1 and v'_2 do define the same points in $\bar{V}_R(G)_1$. This implies that the difference between the number of connected components of $G'(S')$ and $G'[S']$ is equal to $\sharp(V'_R(G)_1) - \sharp(\bar{V}_R(G)_1)$. Those arguments imply

$$g(G'(S')) - g(G'[S']) \equiv \sharp(V'_R(G)_1) - \sharp(\bar{V}_R(G)_1) \pmod{2}. \quad (2.11)$$

Combining this with Formula 2.10 we obtain

$$g(G(S')) \equiv \sharp(V_{\mathbb{R}}(G)_1) - \sharp(\overline{V}_{\mathbb{R}}(G)_1) \pmod{2}. \quad (2.12)$$

By construction $G' \setminus S'$ is a real subgraph of G . Let H be a connected component of $G' \setminus S'$ different from a vertex then $\overline{H}$ is also a connected component of $G' \setminus S'$. Since $G' \setminus S'$ contains no real edge and each real vertex is an isolated component of $G' \setminus S'$ we know H contains no real vertex and no real edge. Since $a(G) = 0$ it follows $H \neq \overline{H}$. But H is isomorphic to $\overline{H}$ hence $g(H) = g(\overline{H})$. This implies $g(G' \setminus S')$ is even, therefore formulas 2.9 and 2.12 imply

$$g(G') \equiv \sharp(V_{\mathbb{R}}(G)_1) - \sharp(\overline{V}_{\mathbb{R}}(G)_1) \pmod{2}. \quad (2.13)$$

Combining this with Formulas 2.4 and 2.8 we obtain the desired formula $g(G) \equiv s(G) + 1 \pmod{2}$.

Step 5. Induction argument on the number of connected components of $G(\mathbb{R})$.

Now we assume $G(\mathbb{R})$ is not connected and fix a connected component G_1 of $G(\mathbb{R})$. Since G is connected there exists another connected component G_2 of $G(\mathbb{R})$ such that there is a walk Γ in G that connects a vertex v_1 of G_1 to a vertex v_2 of G_2 having no real inner vertex. We also can assume that all vertices of Γ are mutually different. Then $\overline{\Gamma}$ satisfies the same properties. Moreover $\overline{\Gamma}$ is different from Γ , indeed, the only edge e of Γ containing v_1 is non-real and $\overline{e}$ is the only edge of $\overline{\Gamma}$ containing v_1 . Let $S = \Gamma \cup \overline{\Gamma}$. Clearly S is a real subgraph of G , hence also $G \setminus E(S)$ is a real subgraph of G . Moreover $(G \setminus E(S))(\mathbb{R}) = G(\mathbb{R})$. Let $G'_1; \dots; G'_t$ be the connected subgraphs of $G \setminus E(S)$. If G'_i contains some real vertex of G then G'_i is a real subgraph of G and $a(G) = 0$ implies $a(G'_i) = 0$. Assume $1 \leq x \leq t$ such that G'_i contains a real vertex of G if and only if $1 \leq i \leq x$. Also assume v_1 is a vertex of G'_1 and in case v_2 is not a vertex of G'_1 , v_2 is a vertex of G'_2 . We are going to assume that the theorem holds for those real subgraphs $G'_1; \dots; G'_x$ and we are going to prove that this implies the theorem holds for G . In case $x \geq 2$ for each $1 \leq i \leq x$ the graph $G'_i(\mathbb{R})$ has less connected components than $G(\mathbb{R})$ and we can use the induction hypothesis. In case $x = 1$, G'_1 is a proper real subgraph of G such that $G'_1(\mathbb{R}) = G(\mathbb{R})$. In particular $G'_1(\mathbb{R})$ is not connected and we can apply the same construction to G'_1 and continuing in this way we arrive at a situation such that $x \geq 2$. Again we can use the induction hypothesis.

For $1 \leq i \leq t$ let $W_i = V(S) \cap V(G'_i)$. Since G is connected $W_i \neq \emptyset$ for $1 \leq i \leq t$ and since $G'_1; \dots; G'_t$ are different connected components of $G \setminus S$ one has $W_i \cap W_j = \emptyset$ for $i \neq j$. Moreover $G(E(S))$ is obtained from $S = G[E(S)]$ by identifying vertices w_1 and w_2 of S if and only if there exists $1 \leq i \leq t$ such that $w_1, w_2 \in W_i$. This implies

$$g(G(E(S))) = g(S) + \left(\sum_{i=1}^t \sharp(W_i) \right) - t. \quad (2.14)$$




Figure 7: Connected components of $G \setminus E(S)$.

We use the decomposition

$$g = g(G) = g(G(E(S))) + g(G \setminus E(S)) = g(G(E(S))) + \sum_{i=1}^t g(G'_i). \quad (2.15)$$

For $1 \leq i \leq x$ we write $s_i = s(G'_i)$ and we assume $s_i \equiv g(G'_i) + 1 \pmod{2}$ and $s_i \leq g(G'_i) + 1$ for $1 \leq i \leq x$. This implies

$$g \geq g(G(E(S))) + \sum_{i=1}^x (s_i - 1) + \sum_{i=x+1}^t g(G'_i). \quad (2.16)$$

with equality if and only if $s_i = g(G_i) + 1$ for $1 \leq i \leq x$. Also by definition $s(G) = \sum_{i=1}^x s_i$.

Let $1 \leq i \leq x$ and assume $w \in W_i \setminus \{v_1; v_2\}$. Since $\overline{G_i} = G_i$ it follows $\overline{w} \in W_i$. In case $v_2 \in V(G'_1)$ it implies $\sharp(W_i)$ is even for $1 \leq i \leq x$ and in case $v_2 \in V(G'_2)$, $\sharp(W_1)$ and $\sharp(W_2)$ are odd and $\sharp(W_i)$ is even for $3 \leq i \leq x$. In both cases we obtain $\sum_{i=1}^x \sharp(W_i)$ is even. In case $v_2 \in V(G'_1)$ it also implies $\sum_{i=1}^x \sharp(W_i) \geq 2x$ and in case $v_2 \in V(G'_2)$ it implies $\sum_{i=1}^x \sharp(W_i) \geq 2x - 2$. Since $g(G[S]) \geq 1$ it follows, combining formulas 2.14 and 2.16 that

$$g(G) \geq 1 + 2x - x + \sum_{i=1}^x (s_i - 1) = s(G) + 1 \quad (2.17)$$

in case $v_2 \in V(G'_1)$ and

$$g(G) \geq 1 + 2x - 2 - x + \sum_{i=1}^x (s_i - 1) = s(G) - 1 \quad (2.18)$$

in case $v_2 \in V(G'_2)$. We obtain $s(G) \leq g(G) + 1$ and moreover the computation implies the following claim we are going to use in the proof of Theorem 3.9.

Claim. If $s(G) = g(G) + 1$ then $v_2 \in V(G_2)$, $\sharp(W_1) = \sharp(W_2) = 1$, $\sharp(W_i) = 2$ for $3 \leq i \leq x$, $\sharp(W_i) = 1$ for $x + 1 \leq i \leq t$, $g(G'_i) = s(G'_i) - 1$ for $1 \leq i \leq x$ (in particular $a(G'_i) = 0$), $g(G'_i) = 0$ for $x + 1 \leq i \leq t$ and $g(S) = 1$ (hence S is a cycle).

In order to finish the proof of the theorem we need to prove $s(G) \equiv g(G) + 1 \pmod{2}$. If S would not be a cycle then there would be an inner vertex w of Γ such that $\bar{w}$ is also a vertex of Γ . The subwalk of Γ that connects w to $\bar{w}$ would imply $a(G) = 0$, hence a contradiction. By induction we have $s_i \equiv g(G'_i) + 1 \pmod{2}$ for $1 \leq i \leq x$. Since G'_i contains no real vertex for $x + 1 \leq i \leq t$ also $\bar{G}'_i \neq G'_i$ in that case. This implies $\sum_{i=x+1}^t g(G'_i) \equiv 0 \pmod{2}$, $t \equiv x \pmod{2}$ and $\sum_{i=x+1}^t \sharp(W_i) \equiv 0 \pmod{2}$. We already noticed that $\sum_{i=1}^x \sharp(W_i) \equiv 0 \pmod{2}$. Therefore, from formulas 2.14 and 2.15 it follows that $g(G) \equiv 1 - x + \sum_{i=1}^x (s_i - 1) \pmod{2}$, hence $g(G) \equiv s(G) + 1 \pmod{2}$. $\square$

Example 2.8. Let g , a and s be nonnegative integers such that $0 \leq s \leq g + 1$ with $s \equiv g + 1 \pmod{2}$ and $a \in \{0, 1\}$ such that $a = 0$ if $s = g + 1$ and $a = 1$ if $s = 0$ (hence g is odd in this case). We give an example of a graph G with a real structure satisfying $g(G) = g$, $a(g) = a$ and $s(G) = s$.

First assume $a = 0$ (hence $s \geq 1$) and write $g + 1 - s = 2x$ (with $x \geq 0$ because $s \leq g + 1$). Take vertices $v_1; \dots; v_s$ and two different edges $e_i; \bar{e}_i$ for $1 \leq i \leq s - 1$ such that $\psi(e_i) = \psi(\bar{e}_i) = \{v_i; v_{i+1}\}$. Then take vertices $w_1; \dots; w_x; \bar{w}_1; \dots; \bar{w}_x$ and edges $f_i; f'_i; \bar{f}_i; \bar{f}'_i$ ($1 \leq i \leq x$) with $\psi(f_1) = \psi(f'_1) = \{v_1; w_1\}$, $\psi(\bar{f}_1) = \psi(\bar{f}'_1) = \{v_1; \bar{w}_1\}$, $\psi(f_i) = \psi(f'_i) = \{w_{i-1}; w_i\}$ and $\psi(\bar{f}_i) = \psi(\bar{f}'_i) = \{\bar{w}_{i-1}; \bar{w}_i\}$. We obtain a graph G such that $g(G) = s - 1 + 2x = g$. Define the real structure on G such that $\bar{\cdot}$ is conjugation and with $\bar{v}_i = v_i$, then $G(\mathbb{R})$ is the disjoint union of the trivial graphs $\{v_1\} \cup \{v_2\} \cup \dots \cup \{v_s\}$, hence $s(G) = s$. Clearly $a(G) = 0$.




Figure 8: Graph with $a = 0$.

Next assume $s = 0$, hence $a = 1$ and g is odd. Write $g + 1 = 2x$ and take two vertices $v; \bar{v}$ and $2x$ edges $e_1; \dots; e_x; \bar{e}_1; \dots; \bar{e}_x$ with $\psi(e_i) = \psi(\bar{e}_i) = \{v; \bar{v}\}$ for $1 \leq i \leq x$. Clearly $g(G) = 2x - 1 = g$. Defining the real structure on G such that $\bar{}$ denotes conjugation we obtain $s(G) = 0$ and clearly $a(G) = 1$.

Finally take $s \neq 0$ but $a = 1$. Write $g + 1 - s = 2x$ with $x \geq 1$ because $s \leq g - 1$. Take s vertices $v_1; \dots; v_s$ and $2s - 2$ edges $e_1; \bar{e}_1; \dots; e_{s-1}; \bar{e}_{s-1}$ with $\psi(e_i) = \psi(\bar{e}_i) = \{v_i; v_{i+1}\}$ for $1 \leq i \leq s - 1$. Take two more vertices v and $\bar{v}$ and $2x + 2$ more edges $f; \bar{f}; f_1; \bar{f}_1; \dots; f_x; \bar{f}_x$ with $\psi(f) = \{v_s; v\}$, $\psi(\bar{f}) = \{v_s; \bar{v}\}$ and $\psi(f_i) = \psi(\bar{f}_i) = \{v; \bar{v}\}$ for $1 \leq i \leq x$. Then $g(G) = s - 1 + 1 + 2x - 1 = g$. Take a real structure on G such that $\bar{}$ denotes conjugation and $\bar{v}_i = v_i$ for $1 \leq i \leq s$. As in the first part of the example we have $s(G) = s$ and since $x \geq 1$, by construction we also have $a(G) = 0$.




Figure 9: Graph with $a = 1$.

3 Real linear systems

We start by recalling some definitions concerning divisors and linear systems on graphs as described in e.g. [1]. A *divisor* on a graph G is a formal integer linear combination of vertices of G . In case D is such a divisor and v is a vertex we write $D(v)$ to denote the coefficient of D at v , hence $D = \sum_{v \in V(G)} D(v)v$. The *support* $\text{Supp}(D)$ of D is the subset of $V(G)$ of vertices v satisfying $D(v) \neq 0$. A divisor is called *effective* if $D(v) \geq 0$ for each vertex $v \in V(G)$ and in that case we write $D \geq 0$. For a divisor D we call $\sum_{v \in V(G)} D(v)$ the degree $\deg(D)$ of D . We write $\text{Div}(G)$ to denote the group of divisors on G (hence it is the free abelian group on $V(G)$). We write $\text{Div}^0(G)$ to denote the subgroup of divisors on G of degree 0.

Let $\mathcal{M}(G)$ be the abelian group of integer-valued functions f on $V(G)$. For $f \in \mathcal{M}(G)$ we define the *principal divisor* $\Delta(f)$ as follows. For $v \in V(G)$ we

define

$$\Delta(f)(v) = \sum_{\substack{e \in E(G) \\ \psi(e) = \{v; w\}}} (f(w) - f(v)). \quad (3.1)$$

Clearly $\Delta(f) \in \text{Div}^0(G)$ and $\{\Delta(f) : f \in \mathcal{M}(G)\}$ is a subgroup $\text{Prin}(G)$ of $\text{Div}^0(G)$. We say two divisors $D_1, D_2 \in \text{Div}(G)$ are linearly equivalent if and only if $D_2 - D_1 \in \text{Prin}(G)$. In particular $\deg(D_1) = \deg(D_2)$ and we write $D_1 \sim D_2$. For a divisor $D \in \text{Div}(G)$ we define the *complete linear system* $|D|$ as follows:

$$|D| = \{D' \in \text{Div}(G) : D' \geq 0 \text{ and } D' \sim D\}$$

We also introduce a notion for the *rank* $\text{rk}(D)$ of D as follows. In case $|D| = \emptyset$ we put $\text{rk}(D) = -1$. So assume $|D| \neq \emptyset$, then $\text{rk}(D)$ is the maximal integer value r such that for each $E \geq 0$ with $\deg(E) = r$ there exists $D' \in |D|$ with $D' - E \geq 0$. For a divisor D on a smooth curve X the corresponding notion is the same as the dimension of the complete linear system $|D|$.

Now assume G has a real structure. Associated to $D = \sum_{v \in V(G)} D(v)v$ there is a conjugated divisor $\overline{D} = \sum_{v \in V(G)} D(\overline{v})v$, hence $\overline{D}(v) = D(\overline{v})$. We say D is a *real divisor* if $D = \overline{D}$. In case $f \in \mathcal{M}(G)$ we define $\overline{f} \in \mathcal{M}(G)$ by $\overline{f}(v) = f(\overline{v})$. Clearly $\Delta(\overline{f}) = \overline{\Delta(f)}$.

Definition 3.1. *Let D be a real divisor on a graph G with a real structure. The real rank $\text{rk}_{\mathbb{R}}(D)$ is the maximal integer value r such that for each real divisor $E \geq 0$ with $\deg(E) = r$ there exists a real divisor $D' \in |D|$ with $D' - E \geq 0$.*

In the case of a real curve X the real dimension of a complete linear system associated to a real divisor D is equal to the dimension of the complete linear system $|D|$ on the complex curve $X_{\mathbb{C}}$ (see e.g. [6, p. 200]). In the graph-case this becomes only one inequality.

Proposition 3.2. *Let G be a graph with a real structure and let D be a real divisor on G . One has $\text{rk}_{\mathbb{R}}(D) \geq \text{rk}(D)$.*

Proof. In case $\text{rk}(D) = -1$ there is nothing to prove, so assume $\text{rk}(D) \geq 0$.

Let E be a real effective divisor of degree r on G . There exists $D'' \in |D|$ such that $D'' - E \geq 0$. Let $f \in \mathcal{M}(G)$ with $D + \Delta(f) = D''$. Since $\overline{D} = D$ we obtain $D + \Delta(\overline{f}) = \overline{D''}$, hence $\overline{D''} \in |D|$.

Let $g = \max\{f; \overline{f}\}$ and let $D' = D + \Delta(g)$. We are going to prove that $D' \geq 0$ (hence $D' \in |D|$); $\overline{D'} = D'$ (hence D' is real) and $D' \geq E$. This will imply the proposition.

Since $D'' \geq 0$, for each $v \in V(G)$ one has

$$D(v) + \sum_{\substack{e \in E(G) \\ \psi(e) = \{v; w\}}} (f(w) - f(v)) = D''(v) \geq 0. \quad (3.2)$$

Using $\bar{v}$ instead of v and taking into account that $D(\bar{v}) = D(v)$, we obtain

$$D(v) + \sum_{\substack{e \in E(G) \\ \psi(e) = \{v; w\}}} (\bar{f}(w) - \bar{f}(v)) = D''(\bar{v}) \geq 0 \quad (3.3)$$

Assume $f(v) \geq f(\bar{v})$, hence $g(v) = f(v)$. For each $e \in E(G)$ with $\psi(e) = \{v; w\}$ one has $g(w) - g(v) \geq f(w) - f(v)$ therefore Formula 3.2 implies $D(v) + \Delta(g)(v) = D'(v) \geq D''(v) \geq 0$. Assume $\bar{f}(v) \geq f(v)$ hence $g(v) = \bar{f}(v)$. Now for e as before we have $g(w) - g(v) \geq \bar{f}(w) - \bar{f}(v)$. Again, using Formula 3.3 we obtain $D(v) + \Delta(g)(v) = D'(v) \geq D''(\bar{v}) \geq 0$. This proves $D' \in |D|$.

By definition $\bar{g} = g$, hence $\Delta(\bar{g}) = \Delta(g)$ and therefore $\bar{D}' = \bar{D} + \Delta(\bar{g}) = D'$. This proves D' is a real divisor.

For each vertex v we have $D''(v) \geq E(v)$. In case v is a real vertex $g(v) = f(v)$ and in such case we obtained $D'(v) \geq D''(v)$ hence $D'(v) \geq E(v)$. Assume v is not a real vertex. Note that $D''(\bar{v}) \geq E(\bar{v}) = E(v)$. We can assume $g(v) = f(v)$ (otherwise we use $\bar{v}$ instead of v) hence $D'(v) \geq D''(v)$ and therefore $D'(v) \geq E(v)$. Since D' and E are both real divisors we also have $D'(\bar{v}) \geq E(\bar{v})$. $\square$

Example 3.3. Let G' be a graph with $g(G') \geq 1$. Fix $v' \in V(G')$. We construct a new graph G by taking two copies G'_1 and G'_2 of G' (hence $v'_i \in V(G'_i)$ is the corresponding copy of v'), a new vertex v and two new edges e_1, e_2 with $\psi(e_i) = \{v; v'_i\}$. On G we define a real structure such that ι induces the identification of G'_1 and G'_2 using their identification with G' , $\bar{e}_1 = e_2$ and v is a real vertex. Then $D = v$ is a real divisor and clearly $\text{rk}_{\mathbb{R}}(D) = 1$. Since $g(G) \neq 0$ there is no linear system of degree 1 and dimension at least 1, hence $\text{rk}(D) = 0$ (see e.g. [3, Lemma 1.1]). This proves that in general Proposition 3.2 cannot have equality.

Lemma 3.4. Assume D_1 and D_2 are real divisors on a graph G with a real structure and $f \in \mathcal{M}(G)$ with $\Delta(f) = D_1 - D_2$, then $\bar{f} = f$

Proof. Since $\Delta(f) = D_2 - D_1$ and D_1 and D_2 are real one has $\overline{\Delta(f)} = \Delta(\bar{f}) = D_2 - D_1$. This implies $\Delta(f - \bar{f}) = 0$. However, for $g \in \mathcal{M}(G)$ the equation $\Delta(g) = 0$ implies g is a constant function on $V(G)$. If not, let $V_0 \subset V(G)$ be the subset of $V(G)$ consisting of all $v \in V(G)$ such that g has minimal value at v . Since $V_0 \neq V(G)$ and G is connected there exists $v_0 \in V_0$, $v \in V(G) \setminus V_0$ and $e \in E(G)$ with $\psi(e) = \{v_0; v\}$. Of course $\Delta(g)(v_0) > 0$ for such vertex v_0 , contradicting $\Delta(g) = 0$.

It follows $f - \bar{f}$ is a constant function on $V(G)$. For each vertex v of $V(G)$ one has $(f - \bar{f})(v) = (f - \bar{f})(\bar{v})$ hence

$$f(v) - \bar{f}(v) = f(\bar{v}) - \bar{f}(\bar{v}) = f(\bar{v}) - f(v)$$

hence $2f(v) = 2f(\bar{v})$. This implies $f(v) = f(\bar{v}) = \bar{f}(v)$ and therefore $f = \bar{f}$. $\square$

In case X is a real curve and C is a connected component of $X(\mathbb{R})$ the parity of the degree of the restriction of real divisors on X to C is invariant

under linear equivalence. This is a basic fact in the study of real linear systems on real curves and in the study of their real projective embeddings. The next theorem shows this also holds for graphs with a real structure.

Theorem 3.5. *Let G be a graph with a real structure and let D_1 and D_2 be two linearly equivalent real divisors on G . Let G' be a connected component of $G(\mathbb{R})$, then $\deg(D_1|_{G'}) \equiv \deg(D_2|_{G'}) \pmod{2}$.*

Proof. Because of Lemma 3.4 there exists $f \in \mathcal{M}(G)$ with $\bar{f} = f$ such that $D_2 = D_1 + \Delta(f)$. Let f' be the restriction of f to $V(G')$. Let $\Delta(f')$ be the associated principal divisor on G' , then $\deg(D_1|_{G'}) = \deg(D_1|_{G'} + \Delta(f'))$. Let $D'_2 = D_1|_{G'} + \Delta(f')$. For $v \in V(G')$ one has

$$D_2(v) = D_1(v) + \sum_{\substack{e \in E(G') \\ \psi(e) = \{v; w\}}} (f'(w) - f'(v)) + \sum_{\substack{e \in E(G) \setminus E(G') \\ \psi(e) = \{v; w\}}} (f(w) - f(v))$$

hence

$$D_2(v) = D'_2(v) + \sum_{\substack{e \in E(G) \setminus E(G') \\ \psi(e) = \{v; w\}}} (f(w) - f(v)).$$

For each $e \in E(G) \setminus E(G')$ with $\psi(e) = \{v; w\}$ one has $\bar{e} \neq e$ and $\psi(\bar{e}) = \{v; \bar{w}\}$ and $f(\bar{w}) - f(v) = \bar{f}(w) - f(v) = f(w) - f(v)$. This proves $\sum_{\substack{e \in E(G) \setminus E(G') \\ \psi(e) = \{v; w\}}} (f(w) - f(v))$ is even, hence $D_2(v) \equiv D'_2(v) \pmod{2}$. So we obtain

$$\deg(D_2|_{G'}) = \sum_{v \in V(G')} D_2(v) \equiv \sum_{v \in V(G')} D'_2(v) \pmod{2}$$

Since $\sum_{v \in V(G')} D'_2(v) = \deg(D'_2) = \deg(D_1|_{G'})$, this finishes the proof. $\square$

In the case of smooth real curves X the parity of the canonical linear system on X (which is a real linear system) is even on each connected component of $X(\mathbb{R})$ (see e.g. [9, Corollary 4.3]). In the case of graphs there is a distinguished canonical divisor

$$K_G = \sum_{v \in V(G)} (\text{val } v - 2)v$$

(here $\text{val } v$ is the number of edges $e \in E(\mathbb{R})$ with $v \in \psi(e)$).

Proposition 3.6. *Let G be a graph with a real structure and let G' be a connected component of $G(\mathbb{R})$, then $\deg(K_G|_{G'})$ is even.*

Proof. On the graph G' there is a canonical divisor

$$K_{G'} = \sum_{v \in V(G')} (\deg_{G'}(v) - 2)v$$

One has $\deg(K'_G) = 2g(G') - 2$, hence it is even. For $v \in V(G')$ the difference $K_G(v) - K_{G'}(v)$ is equal to the number of non-real edges $e \in E(G)$ with $v \in \psi(e)$. Of course this number of edges is even, hence $K_G(v) - K_{G'}(v)$ is even. This proves $\deg(K_G|_{G'})$ is even. $\square$

Definition 3.7. A graph G with a real structure is called an M-graph if G has no isolated real edge and $s(G) = g(G) + 1$.

Remark. If G satisfies $s(G) = g(G) + 1$ but G has isolated real edges then there is an associated M-graph obtained as explained in step 1 of the proof of Theorem 2.7

Definition 3.8. A real divisor D on a graph G with a real structure is called totally real if $\text{Supp}(D) \subset V_{\mathbb{R}}(G)$.

The following result concerning real divisors on M-graphs is a very strong one and the corresponding statement for real divisors on M-curves does not hold.

Theorem 3.9. Let G be an M-graph and let D be an effective real divisor on G then D is linearly equivalent to a totally real effective divisor on G .

Proof. We are going to prove a stronger claim: if $v + \bar{v}$ is a non-real vertex pair on G then there is a real vertex w such that $v + \bar{v} \sim 2w$. The proof is going to make use of notations and claims from the proof of Theorem 2.7.

Of course, in case $G = G(\mathbb{R})$ there is nothing to prove, so we assume $G \neq G(\mathbb{R})$. First assume $G(\mathbb{R})$ is connected. As mentioned in the claim inside step 4 of the proof of Theorem 2.7 we obtained $G(E') = G[V_{\mathbb{R}}(G)] = G(\mathbb{R})$, hence $g(G(E')) = s(G) - 1$, and $g(G') = 0$. In particular each edge $e \in E(G)$ satisfying $\psi(e) \subset V_{\mathbb{R}}(G)$ is a real edge. Let v be a non-real vertex of G . There is a walk Γ that connects v to a real vertex v' such that no inner vertex of Γ is a real vertex. Let $V(v)$ be the set of vertices w of G such that there is a walk with ends v and w and containing no real vertex. Let $T(v)$ be subgraph of the induced subgraph $G[V(v) \cup \{v'\}]$ obtained by omitting the loops at v' . It is a subgraph of G' . Since $g(G') = 0$ the graph $T(v)$ is a tree. Since $G(E') = G[V_{\mathbb{R}}(G)]$ there is no walk that connects a vertex $w \neq v'$ of $T(v)$ to a real vertex different from v' . Since $T(v)$ is a tree there exists a function $f_v \in \mathcal{M}(T(v))$ such that $\Delta(f_v) = v - v'$ on $T(v)$. Let $f \in \mathcal{M}(G)$ such that $f|_{V(T(v))} = f_v$ and $f(v'') = f(v')$ for $v'' \in V(G) \setminus V(T(v))$. It follows that on G we have $\Delta(f) = v - v'$, hence $v \sim v'$. In the same way, using $\overline{T(v)}$ we obtain that $\bar{v} \sim v'$. It follows that $v + \bar{v} \sim 2v'$ on G .

Now assume $G(\mathbb{R})$ has $m > 1$ connected components and assume the theorem holds for M-graphs having less than m connected components. As in step 5 of

the proof of Theorem 2.7 we fix a component G_1 of $G(\mathbb{R})$ and let G_2 be as in that proof. We make use of the results in the claim mentioned in step 5 of the proof of Theorem 2.7. We obtain $G_2 \subset G'_2$, in particular $x \geq 2$, hence $G'_1, \dots, G'_x$ are graphs with a real structure such that $G'_i(\mathbb{R})$ has less than m connected components (say $G'_i(\mathbb{R})$ has m_i connected components and $\sum_{i=1}^x m_i = m$). Also $s(G'_i) = g(G'_i) + 1$ for $1 \leq i \leq x$, hence $G'_1; \dots; G'_x$ are M-graphs and we can apply the induction hypothesis on them.




Figure 10: $v + \bar{v}$ on S .

Let $v + \bar{v}$ be a non-real vertex pair on S (the subgraph used in step 5 of the proof of Theorem 2.7). From the above mentioned claim it follows that S is a cycle. Let Γ be the shortest walk on S from v_1 to v ; let it be $v_1 = w_0 e_0 w_1 e_1 \dots e_{a-2} w_{a-1} e_{a-1} w_a = v$. Let $f(w_i) = f(\bar{w}_i) = i$ for $0 \leq i \leq a$ and $f(w) = a$ for $w \in V(S) \setminus \{w_0; \dots; w_a; \bar{w}_0; \dots; \bar{w}_a\}$. Let $f = 0$ on G'_1 and $f = a$ on G'_2 . For $3 \leq i \leq t$ let $f = f(v_i)$ on G_i with $W_i = \{v_i; \bar{v}_i\}$ in case $i \leq x$ and $W_i = \{v_i\}$ in case $i > x$ (see again the above mentioned claim). Then $v + \bar{v} + \Delta(f) = 2v_1$, hence $v + \bar{v} \sim 2v_1$. In a similar way one finds $v + \bar{v} \sim 2v_2$, hence $2v_1 \sim 2v_2$.

Next assume v is a vertex of some G'_i different from v_i (as above) for some $x+1 \leq i \leq t$. As in the case with $G(\mathbb{R})$ being connected we obtain a tree $T(v)$ contained in G'_i satisfying $T(v) \cap C = \{v_i\}$ and no vertex w of $T(v)$ different from v_i is the end point of an edge not contained in $T(v)$. Since $T(v)$ is a tree there exists $f_v \in \mathcal{M}(T(v))$ with $\Delta(f_v) = v_i - v$ on $T(v)$. Take $f \in \mathcal{M}(G)$ with $f|_{V(T(v))} = f_v$ and $f(v') = f(v_i)$ for $v' \in V(G) \setminus V(T(v))$ then $\Delta(f) = v_i - v$ on G . Using $\bar{f}$ we obtain $\bar{v} \sim \bar{v}_i$ hence $v + \bar{v} \sim v_i + \bar{v}_i$. But $v_i + \bar{v}_i \sim 2v_1$ hence $v + \bar{v} \sim 2v_1$.

Finally assume $v + \bar{v}$ is a non-real vertex pair in some G'_i with $1 \leq i \leq x$. By induction there exists $f_i \in \mathcal{M}(G'_i)$ and a real vertex w of G'_i such that $\Delta(f_i) = 2w - (v + \bar{v})$ on G'_i . From Lemma 3.4 we know that $\bar{f}_i = f_i$, hence




Figure 11: v on G_i with $i > x$.




Figure 12: $v + \bar{v}$ on G_i with $i \leq x$.

$f_i(v_i) = f_i(\bar{v}_i)$ in case $i \geq 3$. Define $f \in \mathcal{M}(G)$ such that $f|_{G'_i} = f_i$ and $f(v') = f_i(v_i)$ for $v' \in V(G) \setminus V(G'_i)$. We obtain $\Delta(f) = 2w - (v + \bar{v})$ on G hence $v + \bar{v} \sim 2w$.

□

Definition 3.10. An M -graph G is called a strong M -graph if $G(\mathbb{R})$ has $g(G)+1$ connected components.

In case G is a strong M -graph it follows that each component G_i of $G(\mathbb{R})$ is a tree. For strong M -graphs one has the following strong result concerning linear systems.

Proposition 3.11. A strong M -graph G has a real linear system g_2^1 (meaning it has a real effective divisor D of degree 2 with $\text{rk}(D) = 1$).

Proof. We already know from Theorem 3.5 that for each non-real vertex pair $v + \bar{v}$ there exists a real vertex w with $v + \bar{v} \sim 2w$. Therefore it is enough to prove the following result: let $v'_1; v'_2$ and $v''_1; v''_2$ be two pairs of real vertices, each pair belonging to one component of $G(\mathbb{R})$, then $v'_1 + v'_2 \sim v''_1 + v''_2$ on G .

First assume $G(\mathbb{R})$ has only one component. Since G is a strong M -graph it follows that G is a tree, hence $v'_1 + v'_2 \sim v''_1 + v''_2$. So assume $G(\mathbb{R})$ has $m \geq 2$

components and assume the proposition holds for strong M-graphs having less than m components. Of course we can assume $v'_1; v'_2$ belong to G_1 . If $v''_1; v''_2$ also belong to G'_1 then we can use the induction hypothesis to G'_1 (in the proof of Theorem 3.5 we found G'_1 is an M-graph and since each connected component of $G'_1(\mathbb{R})$ has genus 0 it is a strong M-graph). There exists $f_1 \in \mathcal{M}(G'_1)$ with $\Delta(f_1) = v'_1 + v'_2 - v''_1 - v''_2$. Define $f \in \mathcal{M}(G)$ such that $f|_{G'_1} = f_1$ and $f(v') = f(v_1)$ for $v' \in V(G) \setminus V(G'_1)$. Then $\Delta(f) = v'_1 + v'_2 - v''_1 - v''_2$ on G . From this part of the proof it follows that $v'_1 + v'_2 \sim 2v_1$. But in the proof of Theorem 3.5 we also found $2v_1 \sim 2v_2$ and $2v_1 \sim v_i + \bar{v}_i$ for $3 \leq i \leq x$. From this part of the proof, if $v''_1; v''_2$ belong to G'_2 (resp. G'_i with $i \geq 3$) we also have $v''_1 + v''_2 \sim 2v_2$ (resp. $v''_1 + v''_2 \sim v_i + \bar{v}_i$) hence $v'_1 + v'_2 \sim 2v_1$ in those cases too, hence $v'_1 + v'_2 \sim v''_1 + v''_2$. $\square$

4 Extensions to metric graphs and tropical curves

In order to study the behavior of the specialization of linear systems on curves to linear systems on graphs as described in [2] with respect to the occurrence of a real structure, we need to extend the concept of graphs with a real structure to the context of metric graphs. As a slightly further generalization we generalize to the context of tropical curves. We also consider generalizations of results on real linear systems on graphs with a real structure to those contexts.

A *weighted graph* G is a graph G having a weight function $w : E(G) \rightarrow \mathbb{R}_{>0}$. For an edge $e \in E(G)$ the number $w(e)$ is called the weight of e .

Definition 4.1. A real structure on a weighted graph G is a real structure on the underlying graph such that for each $e \in E(G)$ one has $w(\bar{e}) = w(e)$.

A *metric graph* Γ is a compact, connected metric space such that each point p on Γ has a neighborhood that is isometric to a star-shaped set of some valence $n_p \geq 1$. A *star-shaped set of valence n* is a metric subspace S of $\mathbb{C}$ obtained as follows. There exists a real number $r > 0$ such that

$$S = \{z \in \mathbb{C} : z = te^{2k\pi i/n} \text{ with } 0 \leq t < r \text{ and } k \in \mathbb{Z}\}$$

A discrete subset V of a metric graph Γ is called a set of vertices of Γ if it contains each point p satisfying $n_p \neq 2$. Once such a set is chosen, we denote it by $V(\Gamma)$ and we call it the set of vertices of Γ . The connected components of $V \setminus V(\Gamma)$ are isometric to line segments, they are called the *edges* of Γ . We write $E(\Gamma)$ to denote the set of edges of Γ . We write $\psi(e)$ to denote the set of the end points of the closure $\bar{e}$ (i.e. $\psi(e) = \bar{e} \cap V(\Gamma)$). The length of an edge e of Γ is denoted by $l(e)$. Since Γ is compact those lengths $l(e)$ are finite real numbers.

Definition 4.2. A real structure on a metric graph Γ is an isometry $\iota : \Gamma \rightarrow \Gamma$ such that ι^2 is the identity.

Let Γ be a metric graph with a real structure. We can assume (and we do) that the set of vertices is invariant under ι . As usual, for $p \in \Gamma$ we write $\bar{p}$ instead of $\iota(p)$ and we say $\bar{p}$ is the *conjugated point*. In case $p = \bar{p}$ we say p is a *real point* on Γ . For $e \in E(G)$ the image $\iota(e)$ is an edge $\bar{e} \in E(G)$. We say e is a *real edge* if e is pointwise fixed by ι . In case $e = \bar{e}$ (e is fixed by ι) either e is real or conjugation on e is given by reflection on e with center the mid-point of e . A real point on Γ is either a real vertex of Γ , a point on a real edge of Γ or a mid-point of a non-real edge e of Γ fixed by ι .

Let G be a weighted graph. Associated to G there is a metric graph $\Gamma(G)$ with bijections $b_V : V(G) \rightarrow V(\Gamma(G))$ and $b_E : E(G) \rightarrow E(\Gamma(G))$ such that for $e \in E(G)$ one has $l(b_E(e)) = w(e)$ and $\psi(b_E(e)) = b_V(\psi(e))$.

Definition/Construction 4.3. *Let G be a weighted graph with a real structure. On the associated metric graph $\Gamma(G)$ we define the associated real structure as follows. We use the same conjugation on $V(\Gamma(G))$ as on $V(G)$ (using b_V). Let $e \in E(G)$ be an edge of G . If e is an non-isolated real edge of G then $\iota|_{b_E(e)}$ is the identity and in case e is an isolated real edge $\iota|_{b_E(e)}$ is the reflection on $b_E(e)$ with center the mid-point of $b_E(e)$. In case $e \in E(G)$ is not real with $\psi(e) = \{v_1; v_2\}$ (of course $v_1 = v_2$ is possible) we use an isometry $b_E(e) \rightarrow b_E(\bar{e})$ mapping $b_V(v_i)$ to $b_V(\bar{v}_i)$. This isometry is unique if $v_1 \neq v_2$ but there are 2 choices if $v_1 = v_2$. In the case when the isometry is not unique the choices are made in a compatible way in order to obtain a real structure.*

Remark. In the specialization described in Section 5 the graph G has no loops, hence the problem of the possible choices in Definition/Construction 4.3 does not occur.

Let Γ be a metric graph. Then there is an associated weighted graph $G(\Gamma)$ obtained by identifying $V(G(\Gamma))$ with $V(\Gamma)$ and $E(G(\Gamma))$ with $E(G)$ and using the same function ψ for $G(\Gamma)$ as for Γ . Moreover for $e \in E(G(\Gamma))$ we put $w(e) = l(e)$. Again a real structure on Γ induces a (now uniquely defined) real structure on $G(\Gamma)$ using the action ι on $V(G)$ and $E(G)$. In case e is a non-real edge of Γ fixed by ι the edge e of $G(\Gamma)$ is an isolated real edge.

Since a weighted graph with a real structure is a graph with a real structure Theorem 2.7 holds in the context of weighted graphs.

On a metric graph Γ with a real structure we write $\Gamma(\mathbb{R})$ to denote the set of real points on Γ . It has a finite number of connected components $\Gamma_1; \dots; \Gamma_{s'}$ each one being a metric subgraph (some of them can be points). Let $s(\Gamma) = \sum_{i=1}^{s'} (g(\Gamma_i) + 1)$. A *path* P in a metric graph Γ is the image of a continuous map $\gamma : [0; 1] \rightarrow \Gamma$ and we say P connects $\gamma(0)$ to $\gamma(1)$. We define $a(\Gamma) = 1$ if there exists a non-real point p on Γ such that p is connected to $\bar{p}$ by a path P not containing any real point of Γ , otherwise we define $a(\Gamma) = 0$. Clearly $g(\Gamma) = g(G(\Gamma))$, $s(\Gamma) = s(G(\Gamma))$ and $a(\Gamma) = a(G(\Gamma))$, hence we obtain the following result on metric graphs with a real structure.

Proposition 4.4. *If Γ is a metric graph with a real structure then $s(\Gamma) \leq g(\Gamma) + 1$; $s(\Gamma) \equiv g(\Gamma) + 1 \pmod{2}$ and $s(\Gamma) \leq g(\Gamma) - 1$ if $a(\Gamma) = 1$.*

Remark. This proposition on metric graphs with a real structure can also be obtained as a special case of so-called Smith Theory in topology. The inequality $s(\Gamma) \leq g(\Gamma) + 1$ follows from [5, Theorem 4.1] while the equality $s(\Gamma) \equiv g(\Gamma) + 1 \pmod{2}$ follows from [5, Theorem 4.3]. From this one obtains the similar statements on graphs in Theorem 2.7. Also the structure on the M-graphs needed to prove Theorem 3.9 and Proposition 3.11 can be deduced from it. The author likes to thank prof. V. Kharlamov for mentioning this relation to Smith Theory.

A *tropical curve* T is a connected metric space being the union of a metric graph Γ and a finite number of unbounded edges e . Such an unbounded edge is isometric to $[0; \infty]$ and satisfies $\overline{T \setminus e} \cap e$ is a unique point on Γ corresponding to 0 and denoted by $e(0)$. Then $\psi(e)$ are the points on e corresponding to 0 and ∞ . We call Γ the *finite part* of T .

Definition 4.5. A real structure on a tropical curve T is an isometry $\iota : T \rightarrow T$ such that ι^2 is the identity.

A real structure on a tropical curve T induces a real structure on the finite part Γ . It also induces an involution on its set of unbounded edges. If e is such an unbounded edge with $e = \bar{e}$ then $e(0) = \bar{e}(0)$ and e is pointwise fixed by ι . In that case e is a real unbounded edge of T . As before a point p on T such that $p = \bar{p}$ is called a *real point* and a real point is either a real vertex of T , a point on a real edge of T or a mid-point on a non-real edge of Γ fixed by ι . The set $T(\mathbb{R})$ has finitely many connected components $T_1; \dots; T_{s'}$ each one being a tropical curve with $T_i \cap \Gamma \neq \emptyset$. As in the case of a metric graph we define numbers $s(T)$ and $a(T)$ and clearly $g(\Gamma) = g(T)$, $s(\Gamma) = s(T)$ and $a(\Gamma) = a(T)$. This implies the following proposition.

Proposition 4.6. Let T be a tropical curve with a real structure, then $s(T) \leq g(T) + 1$, $s(T) \equiv g(T) + 1 \pmod{2}$ and $s(T) \leq g(T) - 1$ if $a(T) = 1$.

Now we extend the results of Section 3 to the context of metric graphs and tropical curves. First we recall the definition of linear systems on metric graphs and tropical curves (see e.g. [8]).

Let Γ be a metric graph or more general a tropical curve. A divisor on Γ is a finite $\mathbb{Z}$ -linear combination D of points on Γ . For a point p on Γ we denote $D(p)$ for the coefficient of D at p . A *rational function* f on Γ is a continuous mapping $f : \Gamma \rightarrow \mathbb{R}$ such that for each edge e of Γ (including the unbounded edges if Γ is a tropical curve) identified isometrically with an interval $I \subset [0; +\infty]$ there is a finite partition $e_1 \cup \dots \cup e_n$ of e in subintervals such that $f|_{e_i}$ is affine with an integer slope. We write $\mathcal{M}(\Gamma)$ to denote the set of rational functions on Γ .

In case $p \in \Gamma$ and $f \in \mathcal{M}(\Gamma)$, $\Delta(f)(p)$ is the sum of slopes of f on Γ in all directions emanating from p . In case p is a vertex of Γ and e is an edge of Γ with $p \in \psi(e)$ there is one such slope associated to e at p denoted by $s_e(f, p)$. In case $p \in \Gamma$ is not a vertex there is a unique edge e of Γ with $p \in e$ and there are two such slopes associated to e at p denoted by $s'_e(f, p)$ and $s''_e(f, p)$. In case f is affine at p , $s'_e(f, p) + s''_e(f, p) = 0$. In this way we define a principal divisor

$\Delta(f) = \sum_{p \in \Gamma} \Delta(f)(p)$ on Γ . Now we can define linear equivalence of divisors on Γ , linear systems on Γ and we can define the rank of a divisor on Γ as we did in the case of graphs.

In case Γ has a real structure, as in the case of graphs for a divisor D on Γ , there is a conjugated divisor $\bar{D}$ and for a rational function f on Γ there is a conjugated rational function $\bar{f}$ on Γ . A divisor D is called a real divisor if $D = \bar{D}$ and it is called totally real if moreover the support of D is contained in $\Gamma(\mathbb{R})$.

A metric graph Γ is called *rational* if the length of each edge of Γ is a rational number. In such case $\Gamma_{\mathbb{Q}}$ is the set of points p of Γ such that there is an edge $e \in E(\Gamma)$ with $p \in e$ and the distance of p to the points of $\psi(e)$ is a rational number. We write $\text{Div}^{\mathbb{Q}}(\Gamma)$ to denote the divisors having support on $\Gamma_{\mathbb{Q}}$. If moreover Γ has a real structure then $\text{Div}_{\mathbb{R}}^{\mathbb{Q}}(\Gamma)$ is the group of real rational divisors on Γ .

Some results on linear systems proved for graphs with a real structure also do hold in the context of metric graphs and tropical curves.

Proposition 4.7. *Let Γ be a metric graph or a tropical curve with a real structure and let D be a real divisor on Γ with $\dim |D| \geq r \geq 1$. Then for each effective real divisor E on Γ with $\deg(E) = r$ there exists a real divisor $D' \in |D|$ with $D' \geq E$.*

Proof. The proof is completely similar to the proof of Proposition 3.2. There exists $f \in \mathcal{M}(\Gamma)$ and $D'' \in |D|$ with $D'' = D + \Delta(f)$ and $D'' \geq E$. Let $g = \max\{f, \bar{f}\}$ and let $D' = D + \Delta(g)$. Since $\bar{g} = g$ and D is real also D' is real. Take $p \in \Gamma$ and let e be an edge of Γ with $p \in e$. If $f(p) \geq \bar{f}(p)$ then $g(p) = f(p)$ hence $s_e(g, p) \geq s_e(f, p)$ in case p is a vertex and $s'_e(g, p) \geq s'_e(f, p)$, $s''_e(g, p) \geq s''_e(f, p)$ in case p is not a vertex. It follows that $D'(p) \geq D''(p)$ in that case. Similarly, in case $\bar{f}(p) \geq f(p)$ we obtain $D'(p) \geq D''(\bar{p})$. This implies D' is effective and $D' \geq E$. □

Lemma 4.8. *Assume D_1 and D_2 are real divisors on a metric graph or a tropical curve Γ with a real structure and assume $f \in \mathcal{M}(\Gamma)$ such that $\Delta(f) = D_2 - D_1$ then $\bar{f} = f$.*

The proof of this lemma is exactly the same as the proof of Lemma 3.4.

Theorem 4.9. *Let Γ be a metric graph or a tropical curve with a real structure and let D_1, D_2 be two linearly equivalent real divisors on Γ . If Γ' is a connected component of $\Gamma(\mathbb{R})$ then $\deg(D_1|_{\Gamma'}) \equiv \deg(D_2|_{\Gamma'}) \pmod{2}$.*

Proof. Let $\Delta(f) = D_1 - D_2$, hence $\bar{f} = f$ by Lemma 4.8. First assume Γ' is a mid-point p of a non-real edge e invariant for ι . Since $\bar{f} = f$ and ι is reflection on e with center p it follows that $s'_e(f, p) = s''_e(f, p)$ hence $\Delta(f)(p)$ is even. Hence $D_1(p) \equiv D_2(p) \pmod{2}$.

Now assume Γ' is a connected component of $\Gamma(\mathbb{R})$ that is not an isolated point unless it is a real vertex of Γ . Use $f' = f|_{\Gamma'}$, similar to f' in the proof of

Theorem 3.5. Take $p \in \Gamma'$. If p is not a vertex of Γ then $\Delta(f')(p) = \Delta(f)(p)$. Since $\deg(D_1|_{\Gamma'} + \Delta(f')) = \deg(D_1|_{\Gamma'})$ such points cannot contradict the claim of the theorem. In case p is a vertex of Γ one can copy the arguments given in Theorem 3.5. □

Definition 4.10. *A metric graph Γ or a tropical curve T with a real structure is called an M-metric graph or an M-tropical curve if $s(\Gamma) = g(\Gamma) + 1$ or $s(T) = g(T) + 1$. It is called a strongly M-metric graph or a strongly M-tropical curve if the number of connected components of $\Gamma(\mathbb{R})$ or $T(\mathbb{R})$ is equal to $g(\Gamma) + 1$ or $g(T) + 1$.*

In this definition there is no need to exclude something similar to isolated real edges as in the case for graphs in Definition 3.7 because they became isolated real points similar to the isolated real vertex as obtained in step 1 of the proof of Theorem 2.7. Since Proposition 4.4 is obtained from Theorem 2.7 in a direct way using weighted graphs, it follows that the structure of M-metric graphs and M-tropical curves is very similar to the structure of M-graphs. Since this structure is the basis in proving Theorem 3.9 and Proposition 3.11 we obtain similar results in the context of metric graphs and tropical curves. We leave the details of the proofs to the reader. It consists of using suited functions on Γ having slope 1 at some part of Γ and being constant on the connected components of the complement.

Theorem 4.11. *Let Γ be an M-metric graph or let T be an M-tropical curve. Each real effective divisor is linearly equivalent to a totally real effective divisor.*

Proposition 4.12. *Let Γ be a strong M-metric graph or let T be a strong M-tropical curve. It has a real linear system g_2^1 .*

5 Specialization of real linear systems on real curves to graphs

In his paper [2] Baker introduces a specialization of linear systems from curves to graphs using a specialization from smooth curves to so-called strongly stable curves. We are going to show that this specialization has good behavior with respect to real structures.

Let T be a smooth geometrically irreducible curve defined over $\mathbb{R}$ and let $o \in T(\mathbb{R})$. Let $\pi : \mathcal{C} \rightarrow T$ be a proper flat morphism of relative dimension 1 such that the general fiber is a smooth geometrically irreducible curve and the total space $\mathcal{C}$ is a smooth surface. The starting point is a local description of this situation. Since T contains a smooth $\mathbb{R}$ -rational point the function field of T is a real field (this means -1 is not a square inside $\mathbb{R}(T)$; see [11, p. 282]). The local ring of T at o is a discrete valuation ring R_o having residue field $\mathbb{R}$. It follows that the completion is a valuation ring R with residue field $\mathbb{R}$ and the quotient field $Q = Q(R)$ again is a real field (see Lemma 5.2). We use the base

extension $\text{Spec}(R) \rightarrow T$ and we obtain an arithmetic surface $\mathfrak{X} \rightarrow R$ (here we write R instead of $\text{Spec}(R)$) having special fiber $X_0 = \pi^{-1}(o)$ (a curve defined over $\mathbb{R}$) and generic fiber X defined over Q . We are going to use some easy facts concerning $R \subset Q$.

Lemma 5.1. *Let Q be a real field containing $\mathbb{R}$ then $Q \otimes_{\mathbb{R}} \mathbb{C}$ is a field.*

Proof. Assume the $\mathbb{R}$ -algebra $Q \otimes_{\mathbb{R}} \mathbb{C}$ is not a field and let M be a maximal ideal of $Q \otimes_{\mathbb{R}} \mathbb{C}$. Since both Q and $\mathbb{C}$ are naturally embedded into $Q \otimes_{\mathbb{R}} \mathbb{C}$ we obtain embeddings $Q \subset Q \otimes_{\mathbb{R}} \mathbb{C}/M$ and $\mathbb{C} \subset Q \otimes_{\mathbb{R}} \mathbb{C}/M$. But $\dim_Q(Q \otimes_{\mathbb{R}} \mathbb{C}) = 2$ hence we obtain $Q \cong Q \otimes_{\mathbb{R}} \mathbb{C}/M$. This would imply $\mathbb{C} \subset Q$ contradicting the fact that Q is a real field. $\square$

Lemma 5.2. *Let R be a complete discrete valuation ring containing $\mathbb{R}$ such that its residue field is equal to $\mathbb{R}$. Then the quotient field is a real field.*

Proof. Assume Q is not a real field. There exists a non-zero element f/g of Q (with $f, g \in R$) such that its square is -1, hence $f^2 + g^2 = 0$ in R . Let v be the valuation of R and let $\pi \in R$ be a uniformizing element (i.e. $v(\pi) = 1$). Let $a = \min\{v(f), v(g)\}$. There exists $f', g' \in \mathbb{R}$ such that $v(f - f'\pi^a)$ and $v(g - g'\pi^a)$ are both at least equal to $a + 1$. At least one of those numbers f', g' is different from 0. From $f^2 + g^2 = 0$ it follows that $f'^2 + g'^2 = 0$. This is impossible, hence a contradiction. $\square$

Lemma 5.3. *Let R be a complete discrete valuation ring containing $\mathbb{R}$ such that its quotient field Q is a real field and its residue field is a finite extension of $\mathbb{R}$. Then the residue field is equal to $\mathbb{R}$.*

Proof. Assume there exists a complete discrete valuation ring R containing $\mathbb{R}$ such that its quotient field is a real field Q and the residue field is isomorphic to $\mathbb{C}$. There exists $x \in R$ such that $v(x^2 + 1) > 0$ (again v is as in the proof of Lemma 5.2). Since Q is a real field we know from Lemma 5.1 that $Q_{\mathbb{C}} = Q \otimes_{\mathbb{R}} \mathbb{C}$ is a field. Let $R_{\mathbb{C}}$ be the complete discrete valuation ring inside $Q_{\mathbb{C}}$ extending R . Since the residue field of R is equal to $\mathbb{C}$ it follows $R_{\mathbb{C}}$ is completely ramified over R . Choose an uniformizing parameter π in $R_{\mathbb{C}}$, then this implies each element of $R_{\mathbb{C}}$ can be written in a unique way as a series $\sum_{n=0}^{\infty} (a_n + b_n x) \pi^n$ with $a_n, b_n \in \mathbb{R}$. Such element belongs to R if and only if $a_n = b_n = 0$ for odd n . Since $i \in R_{\mathbb{C}}$ we have such an expression $i = \sum_{n=0}^{\infty} (a_n + b_n x) \pi^n$. Also $x^2 = -1 + \sum_{j=1}^{\infty} (\alpha_j + \beta_j x) \pi^{2j}$. An expression $-1 = \sum_{n=0}^{\infty} (x_n + y_n x) \pi^n$ implies $x_0 = -1$; $y_0 = 0$ and $x_n = y_n = 0$ for $n \geq 1$.

Now we use

$$i^2 = -1 = \sum_{n=0}^{\infty} (a_n + b_n x)^2 \pi^{2n} + 2 \sum_{\substack{i,j=0 \\ i < j}}^{\infty} (a_i + b_i x)(a_j + b_j x) \pi^{i+j}. \quad (5.1)$$

Writing Equation 5.1 as $-1 = \sum_{n=0}^{\infty} (x_n + y_n x) \pi^n$ we call $x_n + y_n x$ the coefficient of π^n in Equation 5.1.

The only contribution to π^0 in Equation 5.1 comes from the term

$$(a_0 + b_0x)^2 = a_0^2 - b_0^2 + 2a_0b_0x + b_0^2 \sum_{j=1}^{\infty} (\alpha_j + \beta_jx)\pi^{2j}.$$

Hence this term gives no contribution to π^l for some odd l and moreover $a_0^2 - b_0^2 = -1$ and $2a_0b_0 = 0$ implying $a_0 = 0$ and $b_0 = 1$. More generally the terms $(a_n + b_nx)^2\pi^{2n}$ and $(a_i + b_ix)(a_j + b_jx)\pi^{i+j}$ with $i + j$ even do not give contributions to π^l for some odd l .

Assume $k \geq 0$ with $a_i = b_i = 0$ for i odd with $i \leq 2k - 1$. Then the only contribution to π^{2k+1} comes from the term

$$2(a_0 + b_0x)(a_{2k+1} + b_{2k+1}x)\pi^{2k+1} \equiv (-2b_{2k+1} + 2a_{2k+1}x)\pi^{2k+1} \pmod{\pi^{2k+2}}.$$

Since the coefficient of π^{2l+1} in Equation 5.1 has to be equal to 0 one obtains $a_{2k+1} = b_{2k+1} = 0$. This proves $a_l = b_l = 0$ for odd l , hence $i \in R$. This contradicts the fact that Q is a real field. $\square$

Remark. This lemma does not hold in general without assuming completeness. As an example, take the localization of $\mathbb{R}[X]$ at the maximal ideal $\langle X^2 + 1 \rangle$. In this case there exist two extensions of R in $Q_{\mathbb{C}}$.

To define the specialization we put some more conditions on the family $\mathfrak{X} \rightarrow R$. From Lemma 5.1 we know $Q \otimes_{\mathbb{R}} \mathbb{C}$ is a field. Let $R_{\mathbb{C}}$ be the complete valuation ring of $Q_{\mathbb{C}}$ extending R and let $\mathfrak{X}_{\mathbb{C}} \rightarrow R_{\mathbb{C}}$ be obtained from the base change $R \subset R_{\mathbb{C}}$. The closed fiber $X_{0,\mathbb{C}}$ is obtained from X_0 by making the base extension $\mathbb{R} \subset \mathbb{C}$, hence there is a complex conjugation on $X_{0,\mathbb{C}}$. We assume $X_{0,\mathbb{C}}$ is *strongly semistable*. This means we assume $X_{0,\mathbb{C}}$ is reduced, all singular points of $X_{0,\mathbb{C}}$ are nodes and its irreducible components are smooth. Associated to $X_{0,\mathbb{C}}$ there is a dual graph G with a real structure obtained as follows.

Construction 5.4. The set of vertices $V(G)$ corresponds to the set of irreducible components of $X_{0,\mathbb{C}}$. For a vertex $v \in V(G)$ we write C_v for the corresponding component. The set of edges $E(G)$ corresponds to the set of nodes of $X_{0,\mathbb{C}}$. If e is an edge of $E(G)$ then we also write e to denote the node of $X_{0,\mathbb{C}}$ and $\psi(e) = \{v, w\}$ if and only if $e \in C_v \cap C_w$. The complex conjugation on $X_{0,\mathbb{C}}$ induces an involution on both $V(G)$ and $E(G)$ and this defines a real structure on G . As an example, an isolated real edge of G corresponds to a node on $X_{0,\mathbb{C}}$ that is an isolated real point.

We write $\overline{Q}^r$ to denote the real closure of Q and $\overline{Q}$ to denote the closure of Q . Then $\overline{Q} = \overline{Q}^r \otimes_{\mathbb{R}} \mathbb{C}$ (see [11, p. 274]). We write $\overline{X}^r$ (resp. $\overline{X}$) to denote the curve obtained from X by making the base extension $Q \subset \overline{Q}^r$ (resp. $Q \subset \overline{Q}$). On $\overline{Q}$ there is a natural involution extending the complex conjugation on $\mathbb{C}$ and having $\overline{Q}^r$ as its fixed field. We keep calling this complex conjugation and it induces a so-called complex conjugation on $\overline{X}$ too. The image of $P \in \overline{X}$ under




Figure 13: Graph of a real curve

this complex conjugation is again called the conjugated point $\bar{P}$. In case $P = \bar{P}$ we say P is a real point on $\bar{X}$. A divisor on $\bar{X}$ invariant under conjugation is called a real divisor on $\bar{X}$.

Let $Q \subset K \subset \bar{Q}^r$ be an intermediate real field with $[K : Q] = d$ finite. Let R_K be the complete discrete valuation ring extending R having quotient field K . From Lemma 5.3 we know the residue field of R_K is equal to $\mathbb{R}$. Consider the base change $\mathfrak{X} \times_R R_K \rightarrow R_K$, let $R_{K, \mathbb{C}}$ be the complete discrete valuation ring in $K \otimes_R \mathbb{C} = K_{\mathbb{C}}$ and consider the base change $(\mathfrak{X} \times_R R_K)_{\mathbb{C}} \rightarrow R_{K, \mathbb{C}}$. The special fiber is defined over $\mathbb{R}$ and if s is a node then it is an A_{d-1} -singularity of $(\mathfrak{X} \times_R R_K)_{\mathbb{C}}$. The union of those nodes is a 0-dimensional subscheme of $\mathfrak{X} \times_R R_K$ (i.e. it is defined over $\mathbb{R}$). The following lemma implies that the resolution of singularities of all those A_{d-1} -singularities is defined over $\mathbb{R}$.

Lemma 5.5. *Let X be a geometrically irreducible variety defined over $\mathbb{R}$ and let $P \in X(\mathbb{C}) \setminus X(\mathbb{R})$. Let $\bar{X}$ be obtained from X by using a base extension $\mathbb{R} \subset \mathbb{C}$ and let $\tilde{\bar{X}}$ be the blowing-up of $\bar{X}$ at P and $\bar{P}$, then $\tilde{\bar{X}}$ is defined over $\mathbb{R}$*

Proof. We may assume X is affine. Let M_P (resp. $M_{\bar{P}}$) be the maximal ideal corresponding to P (resp. $\bar{P}$) and let $I = M_P \cap M_{\bar{P}}$. Let $I_{\mathbb{R}} = \{f \in I : \bar{f} = f\}$ defining a closed subscheme of X . An element of I can be written as $f_1 + if_2$ with f_1, f_2 defined over $\mathbb{R}$. From $f_1(\bar{P}) + if_2(\bar{P}) = 0$ it follows $f_1(P) - if_2(P) = 0$ hence $f_1(P) = f_2(P) = 0$. This implies $f_1, f_2 \in I_{\mathbb{R}}$. We show that the sheaf of ideals induced by $I_{\mathbb{R}} \otimes_{\mathbb{R}} \mathbb{C}$ on $\bar{X}$ is the sheaf of ideals induced by I . Clearly the stalk of $I_{\mathbb{R}} \otimes_{\mathbb{R}} \mathbb{C}$ at P is included in M_P . In case $f \in I$, writing $f = f_1 + if_2$ as before, we have $f_1 \otimes 1 + f_2 \otimes i \in I_{\mathbb{R}} \otimes_{\mathbb{R}} \mathbb{C}$ maps to f . Assume $f \in M_P$ but $f \notin M_{\bar{P}}$ hence $f(\bar{P}) \neq 0$ and therefore $\bar{f}(P) \neq 0$. Clearly $f\bar{f} \in I_{\mathbb{R}}$. Since $\bar{f}$ is invertible at P we can consider $\bar{f}^{-1}(f\bar{f} \otimes 1)$ mapping to f . Hence the stalk at P is equal to M_P . In a similar way the stalk at $\bar{P}$ is equal to $M_{\bar{P}}$.

Let Q be a point on $\bar{X}$ different from P and $\bar{P}$. If there exists $f \in M_P$ with $f(Q)\bar{f}(Q) \neq 0$ then $f\bar{f} \otimes 1$ maps to an invertible function at Q . Of course the same argument holds using $\bar{P}$ instead of P . Assume for each $f \in M_P \cup M_{\bar{P}}$ one has $f(Q)\bar{f}(Q) = 0$. Since $Q \notin \{P, \bar{P}\}$ there exists $f \in M_P$ and $f' \in M_{\bar{P}}$ with

$f(Q)f'(Q) \neq 0$. Since $\bar{f}(Q)\bar{f}'(Q) = 0$ and $ff' + \bar{f}\bar{f}' \in I_{\mathbb{R}}$ it follows $(ff' + \bar{f}\bar{f}') \otimes 1$ maps to an invertible function at Q . It follows that the stalk at Q is the local ring of X at Q .

Now let $\tilde{X} \rightarrow X$ be the blowing-up of X along $I_{\mathbb{R}}$. This is defined over $\mathbb{R}$. It follows from e.g. [7, B.6.9] that $\tilde{X}$ is a closed subscheme of $\tilde{X} \times_{\mathbb{R}} \mathbb{C}$. But clearly $\tilde{X}$ is invariant under complex conjugation in $\tilde{X} \otimes_{\mathbb{R}} \mathbb{C}$ hence $\tilde{X}$ is defined over $\mathbb{R}$. $\square$

We write $\mathfrak{X}_K$ to denote the resulting smooth arithmetic surface over R_K , it satisfies the same assumptions as $\mathfrak{X} \rightarrow R$. In particular the special fiber of $\mathfrak{X}_{K,\mathbb{C}} \rightarrow R_{K,\mathbb{C}}$ is a strongly semistable curve $X_{0,K,\mathbb{C}}$ defined over $\mathbb{R}$ and associated to it there is a graph G_K with a real structure.

Construction 5.6. Let G_d be the graph obtained from G by subdividing each edge e in d parts. Given a real structure on G we define a real structure on G_d as follows. There is a natural inclusion $V(G) \subset V(G_d)$ and the conjugation on $V(G_d)$ restricts to the known conjugation on $V(G)$. Let $e \in E(G)$ and $\psi(e) = \{v; w\}$ (in general $v = w$ is possible). This edge is replaced by edges $e_1; \dots; e_d$ with $\psi(e_i) = \{v_{i-1}; v_i\}$ satisfying $v_0 = v$ and $v_d = w$ (hence $v_i \notin V(G)$ for $1 \leq i \leq d-1$). First assume $\bar{e} \neq e$ then $\psi(\bar{e}) = \{\bar{v}; \bar{w}\}$ and $\bar{e}$ is replaced by edges $\bar{e}_1; \dots; \bar{e}_d$ with $\psi(\bar{e}_i) = \{\bar{v}_{i-1}; \bar{v}_i\}$ satisfying $\bar{v}_0 = \bar{v}$ and $\bar{v}_d = \bar{w}$. We define $\bar{e}_i = \bar{e}_i$ and $\bar{v}_i = \bar{v}_i$ for $1 \leq i \leq d$. In particular the edges $e_1; \dots; e_d$ and the new vertices $v_1; \dots; v_{d-1}$ are non-real. Assume e is a non-isolated real edge, hence also v and w are real vertices. We define $\bar{v}_i = v_i$ and $\bar{e}_i = e_i$ for $1 \leq i \leq d$. Hence all edges e_i and all new vertices v_i are real. Finally assume e is an isolated real edge. Then $w = \bar{v}$. We define $\bar{v}_i = v_{d-i}$ and $\bar{e}_i = e_{d+1-i}$. Hence all edges e_i and all new vertices v_i are non-real except for vertex $v_{d/2}$ in case d is even and edge $e_{(d+1)/2}$ in case d is odd. In case d is odd $e_{(d+1)/2}$ is an isolated real edge.

Proposition 5.7. *The graph G_K with its real structure is equal to the graph G_d with its real structure.*

Proof. The proper transforms of the components of $X_{0,\mathbb{C}}$ are components of $X_{0,K,\mathbb{C}}$ and they give rise to a natural inclusion $V(G) \subset V(G_K)$. Lemma 5.5 implies that the conjugation on $V(G_K)$ induces the conjugation on $V(G)$. All components corresponding to vertices from $V(G_K) \setminus V(G)$ correspond to new components obtained from resolving the singularities of $(\mathfrak{X} \times_R R_K)_{\mathbb{C}}$. Let $e \in E(G)$ with $\psi(e) = \{v; w\}$. First let e be a non-real edge of G ; it corresponds to a non-real singular point (also denoted by e) of $(\mathfrak{X} \times_R R_K)_{\mathbb{C}}$. Of course, the conjugated point $\bar{e}$ is also a singular point of $(\mathfrak{X} \times_R R_K)_{\mathbb{C}}$. On a small complex neighborhood of e on $(\mathfrak{X} \times_R R_K)_{\mathbb{C}}$ we construct the resolution of the A_{d-1} -singularity obtaining as exceptional divisor a chain of $d-1$ copies of $\mathbb{P}_{\mathbb{C}}^1$ intersecting the proper transforms of C_v and C_w transversally at one point at the ends of the chain (those are the images on C_v and C_w of the singularity e). In $V(G_K)$ this corresponds to subdividing e into d parts. From Lemma 5.5 it follows that using complex conjugation on $\mathfrak{X}_{0,K,\mathbb{C}}$ we obtain the chain of

$\mathbb{P}_{\mathbb{C}}^1$'s obtained from resolving the singularity of $(\mathfrak{X} \times_R R_K)_{\mathbb{C}}$ at $\bar{e}$. On $V(G_K)$ we obtain the subdividing of $\bar{e}$ into d parts and the same conjugation for this part of $V(G_K)$ as on the graph G_d .

Next assume e is a non-isolated real edge of G . It corresponds to a non-isolated real node of $X_{0,\mathbb{C}}$ and locally the equation of $\mathfrak{X} \times_R R_K$ over $\mathbb{R}$ is given by $x^2 - y^2 = t^d$. In this local description $x - y = 0$ and $x + y = 0$ are local equations of C_v and C_w . The blowing-up of $\mathbb{C}^3$ at $(0; 0; 0)$ and the strict transform $\mathfrak{X}'$ of the subset of $(\mathfrak{X} \times_R R_K)_{\mathbb{C}}$ is defined over $\mathbb{R}$. Let $X; Y; T$ be homogeneous coordinates on the exceptional divisor E then the intersection of E with $\mathfrak{X}'$ has equation $X^2 - Y^2 = T^2$ in case $d = 2$ and $X^2 - Y^2 = 0$ in case $d \geq 3$. The intersection points with C_v and C_w are $(1 : 1 : 0)$ and $(1 : -1 : 0)$. In case $d = 2$ we obtain one new component defined over $\mathbb{R}$. This corresponds to a new real vertex v_1 . In case $d \geq 2$ we obtain two new real components corresponding to $X - Y = 0$ and $X + Y = 0$; those correspond to the new vertices v_1 and v_{d-1} . The local equation of $\mathfrak{X}'$ at $(0 : 0 : 1)$ is given by $x^2 + y^2 = t^{d-2}$. Hence if $d = 3$ the resolution is finished, if $d \geq 4$ we have to continue. It follows that in G_K the edge e is divided into d parts and all new edges and vertices are real. This corresponds to the real structure on G_d .

Finally assume e is an isolated real edge. It corresponds to an isolated real node of $X_{0,\mathbb{C}}$ and locally over $\mathbb{R}$ the equation of $(\mathfrak{X} \times_R R_K)_{\mathbb{C}}$ is given by $x^2 + y^2 = t^d$. Using a similar blowing-up the intersection of $\mathfrak{X}'$ and E has equation $X^2 + Y^2 = T^2$ if $d = 2$ and $X^2 + Y^2 = 0$ if $d > 2$. The intersection points with the proper transform C_v and $C_w = C_{\bar{v}}$ are $(1 : i : 0)$ and $(1 : -i : 0)$. Hence they are two conjugated non-real nodes. This proves $\bar{e}_1 = e_d$, hence e_1 (and e_d) are non-real edges. In case $d = 2$ there is a new component v_1 and it is defined over $\mathbb{R}$ (it is the empty conic over $\mathbb{R}$). In case $d \geq 3$ we obtain two conjugated components with equations $X + iY = 0$ and $X - iY = 0$. This shows $\bar{v}_1 = v_{d-1}$. The local equation over $\mathbb{R}$ of $\mathfrak{X}'$ at $(0 : 0 : 1)$ is given by $x^2 + y^2 = t^{d-2}$. In case $d = 3$ the singularity is resolved and the intersection of the new components is an isolated real node corresponding to an isolated real edge on G_K . In case $d \geq 4$ we have to continue. It follows that in G_K the edge e is divided into d parts and the real structure on it corresponds to G_d . $\square$

Now we recall the specialization map studied by Baker in [2]. For a divisor $\mathcal{D}$ on $\mathfrak{X}_{\mathbb{C}}$ and a component C of $X_{0,\mathbb{C}}$ consider the intersection number defined by $(C, \mathcal{D}) = \deg(\mathcal{O}_{\mathfrak{X}}(\mathcal{D})|_C)$. This defines a homomorphism

$$\rho : \text{Div}(\mathfrak{X}) \rightarrow \text{Div}(G) : \mathcal{D} \rightarrow \sum_{v \in V(G)} (C_v, \mathcal{D}) v.$$

Now let D be a divisor on X (hence a divisor on $\bar{X}$ defined over Q). Clearly $D = D_1 - D_2$ with D_1 and D_2 both effective divisors on X , hence closed subschemes of $X \subset \mathfrak{X}$. Take the closures of D_1 and D_2 on $\mathfrak{X}$; their difference is denoted by $cl(D) \in \text{Div}(\mathfrak{X})$. The composition with ρ gives rise to a homomorphism (also denoted by ρ)

$$\rho : \text{Div}(X) \rightarrow \text{Div}(G) : D \rightarrow \rho(\text{cl}(D))$$

Since D is defined over Q the divisor $\text{cl}(D)$ is defined over $\mathbb{R}$ and this implies $\rho(D)$ is a real divisor on G . Let Γ be the metric graph associated with the weighted graph obtained from G giving weight 1 to each edge $e \in E(G)$. The real structure on G induces a real structure on Γ as explained in Definition/Construction 4.3. So we consider Γ as a rational metric graph with a real structure obtained in that way. From the previous map Baker obtains a natural homomorphism $r : \text{Div}(\overline{X}) \rightarrow \text{Div}^{\mathbb{Q}}(\Gamma)$ (see the proof of Theorem 5.8). It is a fundamental observation made by Baker that this map preserves linear equivalence. Now we show that this natural homomorphism also preserves the real structures on X and Γ . The preparations made in this section imply that we can follow directly the arguments from [2].

Theorem 5.8. *If D is a divisor on $\overline{X}^r$ then $r(D)$ is a real divisor on Γ . Hence r induces a natural homomorphism $r : \text{Div}(\overline{X}^r) \rightarrow \text{Div}_{\mathbb{R}}^{\mathbb{Q}}(\Gamma)$.*

Proof. Let D be any divisor on $\overline{X}^r$ (i.e. a real divisor on $\overline{X}$). Let $Q \subset K \subset \overline{K}^r$ be a finite extension such that each real point P and each non-real pair of points $P + \overline{P}$ occurring in $\text{Supp}(D)$ is defined over K . It follows that each point occurring in $\text{Supp}(D)$ as a divisor on $\overline{X}$ is defined over $K_{\mathbb{C}}$. Write $d = [K : Q]$ and consider $\mathfrak{X}_K \rightarrow R_K$. The special fiber $X_{0,K,\mathbb{C}}$ of $\mathfrak{X}_{K,\mathbb{C}} \rightarrow R_{K,\mathbb{C}}$ is defined over $\mathbb{R}$ (this follows from Lemmas 5.3 and 5.5) and from Proposition 5.7 it follows that its dual graph with real structure is equal to G_d with the real structure obtained from G as described in Construction 5.6.

We consider G_d as a weighted graph giving weight $1/d$ to each edge $e \in E(G_d)$. Then the associated metric graph of G_d is equal to Γ and this identification is compatible with the real structures on G_d and Γ . On $\mathfrak{X}_{K,\mathbb{C}}$ each point P of D corresponds to a closed point on the generic fiber, hence its closure is a section $\text{cl}(P)$ of $\mathfrak{X}_{K,\mathbb{C}} \rightarrow R_{K,\mathbb{C}}$. It intersects exactly one component of $X_{0,K,\mathbb{C}}$ corresponding to a vertex $v(P)$ of G_d . This vertex corresponds to rational point $r_K(P)$ on Γ . Then $\rho(D)$ on $\text{Div}(G_K) = \text{Div}(G_d)$ is the sum of those vertices $v(P)$ using P on D , in particular it is a real divisor on G_d . On Γ we obtain $r_K(D) \in \text{Div}_{\mathbb{R}}^{\mathbb{Q}}(\Gamma)$. It is observed by Baker that this divisor on Γ does not depend on the chosen field K , hence it is the divisor $r(D)$ on Γ . $\square$

As a final remark we show that each graph G with a real structure can be obtained from a degeneration $\mathfrak{X} \rightarrow R$. We are going to show that there exists a strongly semistable curve X_0 defined over $\mathbb{R}$ having its dual graph equal to a given graph with a real structure. As a matter of fact we are going to obtain such curve that is totally degenerated (meaning each component of $X_{0,\mathbb{C}}$ is a $\mathbb{P}_{\mathbb{C}}^1$). Then the claim follows from a deep theorem (see e.g. [2, Appendix B]).

Proposition 5.9. *Let G be a graph of genus g without loops having a real structure. There exists a totally degenerated curve X of genus g defined over $\mathbb{R}$ such that the associated graph with real structure is equal to G .*

Proof. A real vertex corresponds to a copy of the real projective line $\mathbb{P}_{\mathbb{R}}^1$ defined over $\mathbb{R}$.

A pair of non-real vertices $v + \bar{v}$ should correspond to a curve Y defined over $\mathbb{R}$ such that $Y(\mathbb{R}) = \emptyset$ and $Y(\mathbb{C})$ is the disjoint union of two copies of the projective line defined over $\mathbb{C}$ and interchanged by complex conjugation. Such a curve is obtained as follows. Let $L, \bar{L}$ be two non-real conjugated lines in $\mathbb{P}_{\mathbb{R}}^2(\mathbb{C})$ and $S = L \cap \bar{L}$, hence $S \in \mathbb{P}_{\mathbb{R}}^2(\mathbb{R})$. The blowing-up of $\mathbb{P}_{\mathbb{R}}^2$ at S is a surface defined over $\mathbb{R}$. The union of the strict transforms of L and $\bar{L}$ is defined over $\mathbb{R}$. This is Y .

Assume $v + \bar{v}$ is a pair of non-real vertices and assume there exist exactly m_1 isolated real edges e and $2m_2$ non-real edges e with $\psi(e) = \{v, \bar{v}\}$. Associated to it we define a scheme Y_{m_1, m_2} defined over $\mathbb{R}$ such that $Y(\mathbb{C})$ has two irreducible components Y_1 and Y_2 both isomorphic to $\mathbb{P}_{\mathbb{C}}^1$, interchanged by complex conjugation and such that $Y_1 \cap Y_2$ consists of nodes of $Y(\mathbb{C})$, m_1 of them being isolated real points and m_2 pairs of conjugated non-real points. In order to obtain this curve we take Y as before, we choose m_1 different pairs of conjugated points $P_1; \bar{P}_1; \dots; P_{m_1}; \bar{P}_{m_1}$ (with $P_i \in Y_1$ and $\bar{P}_i \in Y_2$) and m_2 different pairs of non-conjugated points $Q_{1,1}; Q_{2,1}; \dots; Q_{1,m_2}; Q_{2,m_2}$ with $Q_{i,j} \in Y_i$ and $Q_{2,j} \neq \bar{Q}_{1,j'}$ for all j, j' . Choose an embedding of Y in some $\mathbb{P}^N$ defined over $\mathbb{R}$ such that all chosen points together with their complex conjugated points span a linear subspace Λ of dimension $2m_1 + 4m_2 - 1$ such that Λ intersects Y with multiplicity 1 at those points and contains no other point of Y . Let R_j be a general real point on the line connecting P_j and $\bar{P}_j$ and let R'_j be a general point on the line connecting $Q_{1,j}$ and $Q_{2,j}$. Let Λ' be the linear span of the points R_j and the points R'_j and $\bar{R}'_j$. The projection with center Λ' defines the scheme Y_{m_1, m_2} defined over $\mathbb{R}$.

Now we can construct the curves using suited identifications of points on two schemes Y_1 and Y_2 defined over $\mathbb{R}$ as follows. Let $P_{1,1}; \dots; P_{1,m_1}$ and $P_{2,1}; \dots; P_{2,m_1}$ be different real points on Y_1 and Y_2 . Let $Q_{1,1}; \dots; Q_{1,m_2}$ and $Q_{2,1}; \dots; Q_{2,m_2}$ be different non-real points on Y_1 and Y_2 with $\bar{Q}_{i,j} \neq Q_{i,j}$ (the first index refers to the curve). Using suited embeddings of Y_1 and Y_2 defined over $\mathbb{R}$ and using a Segre embedding of the product one obtains in a similar way as before using a suited projection defined over $\mathbb{R}$ a new curve Y defined over $\mathbb{R}$ obtained from Y_1 and Y_2 by identifying $P_{1,j}$ to $P_{2,j}$; $Q_{1,j}$ to $Q_{2,j}$ and $\bar{Q}_{1,j}$ to $\bar{Q}_{2,j}$. In this way one obtains m_1 new non-isolated real nodes and m_2 new pairs of complex conjugated non-real nodes. They do correspond to m_1 real edges and m_2 pairs of conjugated non-real edges. Iterating this construction one obtains as dual graph the given graph with its real structure. $\square$

References

- [1] M. Baker and S. Norine, *Riemann-Roch and Abel-Jacobi theory on a finite graph*, Adv. Math. **215** (2007), 766-788.
- [2] M. Baker, *Specialization of linear systems from curves to graphs. With an appendix by Brian Conrad*, Algebra Number Theory **2** (2008), 613-788.

- [3] M. Baker and S. Norine, *Harmonic morphisms and hyperelliptic graphs*, Inst. Math. Res. Not. IMRN **15** (2009), 2914-2955.
- [4] J.A. Bondy and U.S.R. Murty, *Graph Theory*, Graduate Texts in Mathematics, vol. 244, Springer-Verlag, 2008.
- [5] G.E. Brendon, *Introduction to compact transformation groups*, Pure and Applied Mathematics, vol. 46, Academic Press, Inc., 1972.
- [6] C. Ciliberto and J. Harris, *Real abelian varieties and real algebraic curves*, Lectures in real geometry (Madrid, 1994), 1996, pp. 167-256.
- [7] W. Fulton, *Intersection theory*, Ergebnisse der Mathematik und ihrer Grenzgebiete, vol. 2, Springer-Verlag, 1984.
- [8] A. Gathmann and M. Kerber, *A Riemann-Roch theorem in tropical geometry*, Math. Z. **259** (2008), 217-230.
- [9] B.H. Gross and J. Harris, *Real algebraic curves*, Ann. Scient. Ec. Norm. Sup. **14** (1981), 157-182.
- [10] J. Huisman, *On the geometry of algebraic curves having many real components*, Revista Matemática Complutense **14** (2001), 83-92.
- [11] S. Lang, *Algebra*, Allison-Wesley Publ. Co., 1965.